

| | | | | | |
|--|--|--|--|---|--|
| FORM PTO-1390 (Modified) (REV. 10-95) | | U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE | | ATTORNEY'S DOCKET NUMBER BC-1003 | |
| TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 | | | | U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR) 09/913655 | |
| | | | | | |
| INTERNATIONAL APPLICATION NO. PCT/US00/09723 | | INTERNATIONAL FILING DATE 12 APRIL 2000 (12.04.00) | | PRIORITY DATE CLAIMED 12 APRIL 1999 (12.04.99) | |
| TITLE OF INVENTION HOMOLOGS OF MAR-BINDING FILAMENT-LIKE PROTEIN 1 (MFP1) | | | | | |
| APPLICANT(S) FOR DO/EO/US HARDER, Patricia A. et al. | | | | | |
| <p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information</p> <ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input checked="" type="checkbox"/> This is an express request to being national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b)) and PCT Articles 22 and 39(1). <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <input checked="" type="checkbox"/> A copy of the International Application was filed (35 U.S.C. 371 (c) (2)) <ol style="list-style-type: none"> <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US) <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371 (c) (2)). <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c) (3)) <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)). <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). <input type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409) <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)). <p>Items 13 to 18 below concern document(s) or information included :</p> <ol style="list-style-type: none"> <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <input type="checkbox"/> A FIRST preliminary amendment. A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input checked="" type="checkbox"/> A change of power of attorney and/or address letter. <input checked="" type="checkbox"/> Certificate of Mailing by Express Mail. <input type="checkbox"/> Other items or information: <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> <p>17. General Power of Attorney 18. Express Mailing Label No.: EJ376014776US</p> </div> | | | | | |

ATTORNEY'S DOCKET NUMBER

BC-1003

20. The following fees are submitted

CALCULATIONS PTO USE ONLY

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) – (5)) :

- | | | |
|-------------------------------------|---|-----------|
| <input checked="" type="checkbox"/> | Search Report has been prepared by the EPO or JPO | \$860.00 |
| <input type="checkbox"/> | International preliminary examination fee paid to USPTO (37 CFR 1.482) | \$690.00 |
| <input type="checkbox"/> | No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) | \$760.00 |
| <input type="checkbox"/> | Neither international preliminary examination fee paid to USPTO (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO | \$1000.00 |
| <input type="checkbox"/> | International preliminary examination fee paid to USPTO (37 CFR 1.482) And all claims satisfied provisions of PCT Article 33(2)-(4) | \$ 100.00 |

ENTER APPROPRIATE BASIC FEE AMOUNT

\$860.00

Surcharge of **\$130.00** for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)).

☐ 20 ☐ 30

\$0.00

| CLAIMS | NUMBER FILED | NUMBER EXTRA | RATE |
|--------------------|--------------|--------------|-------|
| Total Claims | 37 - 20 = | 17 x | \$180 |
| Independent Claims | 4 - 3 = | 17 x | \$80 |

\$306.00

\$80.00

Multiple Dependent Claims (check if applicable)

☒

\$270.00

TOTAL OF ABOVE CALCULATIONS

\$656.00

Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (check if applicable).

\$0.00

SUBTOTAL

\$656.00

Processing Fee of **\$130.00** for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)).

☐ 20 ☐ 30

\$0.00

TOTAL NATIONAL FEE

\$1,516.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) **(check if applicable)**.

☐

\$0.00

TOTAL FEES ENCLOSED

\$1,516.00

Amount to be :
refunded

\$

Charged

\$

- ☐ A check in the amount of _____ to cover the above fees enclosed.
- ☒ Please charge my Deposit Account No. **04-1928** in the amount of **\$1,516.00** to cover the above fees.
- ☒ The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. **04-1928** a duplicate copy of this sheet is enclosed.

NOTE : Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (CFR 1.37(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

**FELTHAM, Neil S.
E. I. DU PONT DE NEMOURS AND COMPANY
Legal Patent Records Center
1007 Market Street
Wilmington, Delaware 19898
United States of America**

SIGNATURE

FELTHAM, NEIL S.

NAME _____

36.506

REGISTRATION NUMBER

DATE _____

8. AUG. 2001 15:43

EPA MUENCHEN +49 89 23994465

NR. 7913 S. 12/19

30-10-2000

00920238.3(12-04-2000) - US00/09723(12-04-2000)

P. CLMSPAMD

14 30 10 00

INTERNATIONAL PRELIMINARY EXAMINATION AUTHORITY

In the Application of:

E.I. DUPONT DE NEMOURS AND COMPANY

Case No: BC1003 PCT

International Application No.: PCT/US00/09723

International Filing Date: 12/04/2000

For: HOMOLOGS OF MAR-BINDING FILAMENT-LIKE PROTEIN 1 (MFP1)

Wilmington Delaware, USA

September 26, 2000

AMENDMENT UNDER ARTICLE 34

The European Patent Office
Erhardtstrasse 27
D-80298 Munich 2, Germany

Sir:

Please cancel Claims 1-20 and substitute the following fifty-six (56) claims:

1. An isolated nucleic acid fragment encoding a tobacco MFP1 protein selected from the group consisting of:
 - (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence selected from the group consisting of SEQ ID NO:2, and SEQ ID NO:4;
 - (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence selected from the group consisting of SEQ ID NO:2, and SEQ ID NO:4;

Printed:02-08-2001

AMENDED SHEET

1

30-10-2000

00920238.3(12-04-2000) - US00/09723(12-04-2000)

POLMSFAMD

11 30 10 00

- (c) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence of (a) or (b) under the following hybridization conditions: 5 x Denhardts, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C;
 - (d) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:14, and SEQ ID NO:15, under the following hybridization conditions: 5 x Denhardts, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C; and
 - (e) an isolated nucleic acid fragment that is complementary to (a), (b), (c) or (d).
2. The isolated nucleic acid fragment of Claim 1 selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:14, and SEQ ID NO:15.
 3. A polypeptide encoded by the isolated nucleic acid fragment of Claim 1.
 4. The polypeptide of Claim 3 selected from the group consisting of SEQ ID NO:2, and SEQ ID NO:4.
 5. An isolated nucleic acid fragment encoding a tobacco MFP1 polypeptide, the peptide having at least 77% identity to SEQ ID NO:17.
 6. An MFP1 polypeptide encoded by the nucleic acid fragment of Claim 5.
 7. A chimeric gene comprising the isolated nucleic acid fragment of either of Claims 1 or 5 operably linked to suitable regulatory sequences.
 8. A transformed host cell comprising a host cell and the chimeric gene of Claim 7.
 9. The transformed host cell of Claim 8 wherein the host cell is a plant cell.
 10. The transformed host cell of Claim 8 wherein the host cell is *E. coli*.
 11. A method of altering the level of expression of a plant MFP1 protein in a host cell comprising:
 - (a) transforming a host cell with the chimeric gene of Claim 7 and;
 - (b) growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric generesulting in production of altered levels of a plant MFP1 protein in the transformed host cell relative to expression levels of an untransformed host cell.
 12. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a tobacco MFP1 protein comprising:
 - (a) probing a cDNA or genomic library with the nucleic acid fragment of Claim 1;

30-10-00

- (b) identifying a DNA clone that hybridizes with the nucleic acid fragment of Claim 1; and
- (c) sequencing the cDNA or genomic fragment that comprises the clone identified in step (b),

wherein the sequenced cDNA or genomic fragment encodes a tobacco MFP1 protein

13. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a tobacco MFP1 protein comprising:

- (a) synthesizing at least one oligonucleotide primer corresponding to a portion of the sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:14, and SEQ ID NO:15;
- (b) amplifying a cDNA insert present in a cloning vector using the oligonucleotide primer of step (a);

wherein the amplified cDNA insert encodes a tobacco MFP1 protein.

14. The product of the method of Claims 12 or 13.

15. An isolated nucleic acid fragment encoding a soybean MFP1 protein selected from the group consisting of:

- (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence as set forth in SEQ ID NO:20;
- (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence as set forth in SEQ ID NO:20;
- (c) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence of (a) or (b) under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C;
- (d) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence as set forth in SEQ ID NO:19 under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C; and
- (e) an isolated nucleic acid fragment that is complementary to (a), (b), (c) or (d).

16. The isolated nucleic acid fragment of Claim 15 as set forth in SEQ ID NO:19.

17. A polypeptide encoded by the isolated nucleic acid fragment of Claim 15.

18. The polypeptide of Claim 17 as set forth in SEQ ID NO:20.

11 30.10.00

19. A nucleic acid fragment, isolated from soybean, encoding an MFP1 polypeptide, the polypeptide having at least 46% identity to SEQ ID NO:17 over a length of 388 amino acids as compared by the Jotun-Hein algorithm.

20. An MFP1 polypeptide encoded by the nucleic acid fragment of Claim 19.

21. A chimeric gene comprising the isolated nucleic acid fragment of either of Claims 15 or 19 operably linked to suitable regulatory sequences.

22. A transformed host cell comprising a host cell and the chimeric gene of Claim 21.

23. The transformed host cell of Claim 22 wherein the host cell is a plant cell.

24. The transformed host cell of Claim 22 wherein the host cell is *E. coli*.

25. A method of altering the level of expression of a plant MFP1 protein in a host cell comprising:

- (a) transforming a host cell with the chimeric gene of Claim 21 and;
- (b) growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric gene

resulting in production of altered levels of a plant MFP1 protein in the transformed host cell relative to expression levels of an untransformed host cell.

26. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a soybean MFP1 protein comprising:

- (a) probing a cDNA or genomic library with the nucleic acid fragment of Claim 15;
- (b) identifying a DNA clone that hybridizes with the nucleic acid fragment of Claim 15; and
- (c) sequencing the cDNA or genomic fragment that comprises the clone identified in step (b),

wherein the sequenced cDNA or genomic fragment encodes a soybean MFP1 protein

27. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a soybean MFP1 protein comprising:

- (a) synthesizing at least one oligonucleotide primer corresponding to a portion of the sequence as set forth in SEQ ID NO:19;
- (b) amplifying a cDNA insert present in a cloning vector using the oligonucleotide primer of step (a);

wherein the amplified cDNA insert encodes a corn MFP1 protein.

28. The product of the method of Claims 26 or 27.

29. An isolated nucleic acid fragment encoding a corn MFP1 protein selected from the group consisting of:

11 30 10 00

- (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence as set forth in SEQ ID NO:22;
 - (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence as set forth in SEQ ID NO:22;
 - (c) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence of (a) or (b) under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C;
 - (d) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence as set forth in SEQ ID NO:21 under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C; and
 - (e) an isolated nucleic acid fragment that is complementary to (a), (b), (c) or (d).
30. The isolated nucleic acid fragment of Claim 29 as set forth in SEQ ID NO:21.
31. A polypeptide encoded by the isolated nucleic acid fragment of Claim 29.
32. The polypeptide of Claim 31 as set forth in SEQ ID NO:22.
33. A nucleic acid fragment, isolated from corn, encoding an MFP1 polypeptide, the polypeptide having at least 40% identity to SEQ ID NO:17, over a length of about 672 amino acids as compared by the Jotun-Hein algorithm.
34. An MFP1 polypeptide encoded by the nucleic acid fragment of Claim 33.
35. A chimeric gene comprising the isolated nucleic acid fragment of either of Claims 29 or 33 operably linked to suitable regulatory sequences.
36. A transformed host cell comprising a host cell and the chimeric gene of Claim 35.
37. The transformed host cell of Claim 36 wherein the host cell is a plant cell.
38. The transformed host cell of Claim 36 wherein the host cell is *E. coli*.
39. A method of altering the level of expression of a plant MFP1 protein in a host cell comprising:
- (a) transforming a host cell with the chimeric gene of Claim 35 and;
 - (b) growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric gene
- resulting in production of altered levels of a plant MFP1 protein in the transformed host cell relative to expression levels of an untransformed host cell.

11 30 10 00

40. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a corn MFP1 protein comprising:

- (a) probing a cDNA or genomic library with the nucleic acid fragment of Claim 29;
- (b) identifying a DNA clone that hybridizes with the nucleic acid fragment of Claim 29; and
- (c) sequencing the cDNA or genomic fragment that comprises the clone identified in step (b),

wherein the sequenced cDNA or genomic fragment encodes a corn MFP1 protein

41. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a corn MFP1 protein comprising:

- (a) synthesizing at least one oligonucleotide primer corresponding to a portion of the sequence as set forth in SEQ ID NO:21;
- (b) amplifying a cDNA insert present in a cloning vector using the oligonucleotide primer of step (a);

wherein the amplified cDNA insert encodes a corn MFP1 protein.

42. The product of the method of Claims 40 or 41.

43. An isolated nucleic acid fragment encoding a rice MFP1 protein selected from the group consisting of:

- (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence as set forth in SEQ ID NO:24;
- (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence as set forth in SEQ ID NO:24;
- (c) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence of (a) or (b) under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C;
- (d) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence as set forth in SEQ ID NO:23 under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C; and
- (e) an isolated nucleic acid fragment that is complementary to (a), (b), (c) or (d).

44. The isolated nucleic acid fragment of Claim 43 as set forth in SEQ ID NO:23.

45. A polypeptide encoded by the isolated nucleic acid fragment of Claim 43.

30-10-2000

00920238.3(12-04-2000) US00/09723(12-04-2000)

P. 01/18/19

M 30.10.00

46. The polypeptide of Claim 45 as set forth in SEQ ID NO:24.
47. A nucleic acid fragment, isolated, from rice, encoding a rice MFP1 polypeptide, the polypeptide having at least 39% identity to SEQ ID NO:17 over a length of 107 amino acids as compared by the Jotun-Hein algorithm.
48. An MFP1 polypeptide encoded by the nucleic acid fragment of Claim 47.
49. A chimeric gene comprising the isolated nucleic acid fragment of either of Claims 43 or 47 operably linked to suitable regulatory sequences.
50. A transformed host cell comprising a host cell and the chimeric gene of Claim 49.
51. The transformed host cell of Claim 50 wherein the host cell is a plant cell.
52. The transformed host cell of Claim 50 wherein the host cell is *E. coli*.
53. A method of altering the level of expression of a plant MFP1 protein in a host cell comprising:
- (a) transforming a host cell with the chimeric gene of Claim 43 and;
 - (b) growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric gene
- resulting in production of altered levels of a plant MFP1 protein in the transformed host cell relative to expression levels of an untransformed host cell.
54. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a rice MFP1 protein comprising:
- (a) probing a cDNA or genomic library with the nucleic acid fragment of Claim 43;
 - (b) identifying a DNA clone that hybridizes with the nucleic acid fragment of Claim 43; and
 - (c) sequencing the cDNA or genomic fragment that comprises the clone identified in step (b),
- wherein the sequenced cDNA or genomic fragment encodes a rice MFP1 protein
55. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a rice MFP1 protein comprising:
- (a) synthesizing at least one oligonucleotide primer corresponding to a portion of the sequence as set forth in SEQ ID NO:23;
 - (b) amplifying a cDNA insert present in a cloning vector using the oligonucleotide primer of step (a);
- wherein the amplified cDNA insert encodes a rice MFP1 protein.
56. The product of the method of Claims 54 or 55.

8. AUG. 2001 15:45
30-10-2000

EPA MUENCHEN +49 89 23994465

NR. 7913 S. 19/19

00920238.3(12-04-2000) - US00/09723(12-04-2000)


P. 01MSPAMD

11 30.10.00

DISCUSSION

By virtue of the these amendments Applicant's invention is drawn to nucleic acids and polypeptides corresponding to Tobacco MFP1 (Claims 1-14), Soybean MFP1 (Claims 15-28), Corn MFP1 (Claims 29-42) and Rice MFP1 (Claims 43-56), and genetic chimera and transformants containing the same, and their methods of use.

Very truly yours,



S. NEIL FELTHAM
Attorney for Applicant
Registration No.: 36,506
Telephone: (302) 992-6460
Fax No.: (302) 892-7949

Printed:02-08-2001

AMENDED SHEET

8

4/pst

TITLE
HOMOLOGS OF MAR-BINDING
FILAMENT-LIKE PROTEIN 1 (MFP1)

This application claims the benefit of U.S. Provisional Application
5 No. 60/128,900, filed April 12, 1999.

FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. This invention
pertains to nucleic acid fragments encoding proteins that are homologs to the
MAR-binding filament-like protein 1 (MFP1) from tomato. More specifically,
10 this invention pertains to two tobacco MFP1 genes and MFP1 homologs from
corn, soybean and rice.

BACKGROUND OF THE INVENTION

The nuclear matrix hypothesis proposes a structural framework for the
eukaryotic nucleus that is similar to the cytoskeleton. To date, its best
15 characterized component is the lamina, a filamentous protein network that lines
the inner membrane of the nuclear envelope. Major components of the lamina
include a group of intermediate-filament (IF) proteins, collectively known as
nuclear lamins, that are classified as type A, B, and C (McKeon *et al.*, *Nature*
319:463-468 (1986)). Lamin B is attached to the inner nuclear membrane via a
20 C-terminal C15 farnesyl group (Schafer *et al.*, *Annu. Rev. Genet.* 30:209-237
(1992)), whereas lamins A and C bind to lamin B. Other integral membrane
proteins interact with lamin B and most likely stabilize the membrane attachment
of lamins (Furukawa *et al.*, *EMBO J.* 14:1626-1636 (1995)). Recent studies have
also demonstrated the ability of lamins A and B to bind DNA, suggesting a role
25 for mammalian lamins in anchoring chromatin to the nuclear envelope. The
interaction between nuclear envelope, lamina, and chromatin is considered to be
of fundamental importance for higher order chromosome organization, as well as
the assembly and disassembly of the nuclear envelope during mitosis (Furukawa
et al., *EMBO J.* 14:1626-1636 (1995)).

30 The nuclear matrix is a second structural skeleton that has been
biochemically defined as the insoluble component that remains after treatment of
isolated nuclei with DNase I and extraction of proteins with high-salt solutions
(Berezney *et al.*, *Biochem. Biophys. Res. Comm.* 60:1410-1417 (1974)) or the
chaotropic agent lithium diiodosalicylate (Mirkowitch *et al.*, *Cell* 39:223-232
35 (1984)). Chromatin binds to the nuclear matrix via matrix attachment regions
(MARs) in the DNA. MARs are generally AT-rich DNA sequences that are
several hundred base pairs long and localized to noncoding regions of the DNA,
but often flanking genes (Gasser *et al.*, *Trends Genet.* 3:16-22 (1987)). However,

there is no consensus sequence known for MARs. The significance of structural characteristics for MARs such as DNA bending and a narrow minor groove due to oligo(dA) tracts has been previously proposed. MARs have been shown to increase transcriptional activity of a linked gene and to confer position-
5 independent, copy-number dependent expression in stably transfected cells (Phi-Wan et al., *EMBO J.* 7:655-664 (1988)).

A small number of MAR binding proteins have been identified from animal nuclei, and they are considered to be components of the nuclear matrix (von Kries et al., *Cell* 64:123-135 (1991); Dickinson et al., *Cell* 70:631-645
10 (1992); Romig et al., *EMBO J.* 11:3431-3440 (1992); Tsutsui et al., *J. Biol. Chem.* 268:12886-12894 (1993); Renz et al., *Nucleic Acids Res.* 24:843-849 (1996); U.S. 5,652,340). In addition, it has been shown that lamins specifically bind to MARs (Luderus et al., *Mol. Cell. Biol.* 14:6297-6305 (1994)). The specific interaction
15 between DNA and the nuclear matrix/nuclear lamina is most likely an important mechanism for long-range gene regulation and higher order chromatin organization (Gasser et al., *Trends Genet.* 3:16-22 (1987)).

Most investigations into structural components of the nucleus have focused on proteins in vertebrates and *Drosophila*, but even in these organisms, our knowledge about the molecular constituents of the nuclear matrix is sparse.
20 Significantly less information is available for other eukaryotes, and in particular for plants. Proteins that are immunologically related to animal IF proteins and lamins have been detected in pea and carrot nuclei (Beven et al., *J. Cell Sci.* (1991) 98 (3), 293-30; McNulty et al., *J. Cell Sci.* 103:407-414 (1992)). Plant nuclear matrix preparations that bind to animal MARs have been reported,
25 suggesting that proteins with similar DNA binding specificities exist in plants as well (Hall et al., *Proc. Natl. Acad. Sci. USA* 88:9320-9324 (1991)).

Effects of MARs on gene expression in plants have been reported, but have been quite variable. In some experimental systems, no reduction of variability but an increase in expression level has been reported (Breyne et al.,
30 *Plant Cell* 4:463-471 (1992); Allen et al., *Plant Cell* 5:603-613 (1993); Allen et al., *Plant Cell* 8:899-913 (1996); U.S. 5,773,689). Other authors have found no significant increase in expression level, but a reduction of variability (van der Geest et al., *Plant J.* 6:413-423 (1994); Mlynarova et al., *Plant Cell* 6:417-426 (1994)). It is not clear what causes these observed differences, but they
35 will most probably be due to the fact that MARs establish different molecular interactions, which might either depend on the features of the MAR itself or on the specific molecular environment of the transformed cell/tissue. The routine use of MARs for strategies to improve transgene expression will greatly depend on the

characterization of the proteins involved in DNA-nuclear matrix attachment and the factors responsible for the observed increase in gene expression.

Currently, no sequence information is available for plant lamin-like proteins. However, the cloning of the cDNA for a plant MAR-binding protein, MFP1, from tomato has been reported (Meier et al., *Plant Cell* 8:2105-2115 (1996)). MFP1 has structural features of a filament-like protein and it preferentially binds to MAR DNA sequences from both plants and animals. In contrast to other known MAR binding proteins, MFP1 contains a hydrophobic N-terminal amino acid sequence that might function as a membrane-spanning domain. MFP1, therefore, has features of a novel anchor protein that most likely connects chromatin via MAR DNA with the nuclear envelope and nuclear filament proteins.

In order to routinely use the attachment of transgenes to the nuclear matrix improve gene expression, it will be necessary to further characterize the elements involved in this process and to better understand the underlying mechanisms. Thus, a need exists to identify and characterize additional nuclear matrix proteins. The present invention presents MFP1-like proteins from other plant species. Furthermore, the present invention shows that a single, immunologically related protein of comparable size is present in a variety of higher-plant species, including important crop plants. This invention pertains to the isolation of cDNAs corresponding to two tobacco MFP1 genes and the characterization of the MFP1 gene family in tobacco. The invention also pertains to the identification and partial characterization of EST sequences from corn, soybean and rice encoding MFP1 proteins from these crop species.

SUMMARY OF THE INVENTION

The present invention provides an isolated nucleic acid fragment encoding a plant MFP1 protein selected from the group consisting of: (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:20, SEQ ID NO:22 and SEQ ID NO:24; (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:20, SEQ ID NO:22 and SEQ ID NO:24; (c) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence of (a) or (b) under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C; (d) an isolated nucleic acid molecule that hybridizes with a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3,

SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:19, SEQ ID NO:21 and SEQ ID NO:23 under the following hybridization conditions: 5 x Denhards, 5 x SSPE, 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C; and (e) an isolated nucleic acid fragment that is
5 complementary to (a), (b), (c) or (d).

Additionally the invention provides a nucleic acid fragment, isolated from corn, encoding an MFP1 polypeptide, the polypeptide having at least 40% identity to SEQ ID NO:17, over a length of about 672 amino acids as compared by the Jotun-Hein algorithm.

10 Similarly the invention provides a nucleic acid fragment, isolated from soybean, encoding an MFP1 polypeptide, the polypeptide having at least 46% identity to SEQ ID NO:17 over a length of 388 amino acids as compared by the Jotun-Hein algorithm.

In another embodiment the invention provide a nucleic acid fragment,
15 isolated, from rice, encoding an MFP1 polypeptide, the polypeptide having at least 39% identity to SEQ ID NO:17 over a length of 107 amino acids as compared by the Jotun-Hein algorithm.

In an alternate embodiment the invention provides an isolated nucleic acid fragment encoding a plant MFP1 polypeptide, the peptide having at least 77%
20 identity to SEQ ID NO:17.

The invention further provides polypeptides encoded by the isolated nucleic acid fragments of the present invention.

In another embodiment the invention provides a chimeric gene comprising the isolated nucleic acid fragment of the present invention operably linked to
25 suitable regulatory sequences.

The invention additionally provides a method of altering the level of expression of a plant MFP1 protein in a host cell comprising: (a) transforming a host cell with the chimeric gene of the present invention and; (b) growing the transformed host cell produced in step (a) under conditions that are suitable for
30 expression of the chimeric gene resulting in production of altered levels of a plant MFP1 protein in the transformed host cell relative to expression levels of an untransformed host cell.

The invention additionally provides transformed host cells comprising the chimeric genes of the present invention.

35 In an alternate embodiment the invention provides methods of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a plant MFP1 protein using portions of the present nucleic acid sequences as hybridization probes or as primers.

BRIEF DESCRIPTION OF THE DRAWINGS,
AND SEQUENCE DESCRIPTIONS

Figure 1 shows a schematic representation of the subfragments E-196 and H-207 that were expressed in *Escherichia coli*.

5 Figure 2A is a gel showing the immunological identification of MFP1-like proteins in different plant species using the aR50 antibody raised against a Le MFP1 polypeptide.

Figure 2B is a gel showing the immunological identification of MFP1-like proteins in different plant species using the a288 antibody raised against a Le
10 MFP1 polypeptide.

Figure 3 shows the schematic structure of the partial cDNAs isolated from a *Nicotiana tabacum* lambda ZAP cDNA library.

Figure 4A shows the percent identical amino acids in pairwise comparisons of the four MFP1 proteins.

15 Figure 4B shows the hydrophilicity and secondary structure analysis of LeMFP1, NtMFP1-1 and AtMFP1.

Figure 5 shows the genomic organization of tobacco MFP1.

The invention can be more fully understood from the following detailed description and the accompanying sequence descriptions which form part of this
20 application.

The following sequence descriptions and sequence listings attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825. The Sequence Descriptions contain the one letter code for nucleotide sequence
25 characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IYUB standards described in *Nucleic Acids Research* 13:3021-3030 (1985) and in the *Biochemical Journal* 219(2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

30 SEQ ID NO:1 is the nucleotide sequence for NtMFP1-1.

SEQ ID NO:2 is the deduced amino acid sequence for NtMFP1-1, encoded by SEQ ID NO:1.

SEQ ID NO:3 is the nucleotide sequence for NtMFP1-2.

35 SEQ ID NO:4 is the deduced amino acid sequence for NtMFP1-2, encoded by SEQ ID NO:3.

SEQ ID NO:5 is the nucleotide sequence which codes for E-196 polypeptide fragment isolated from tomato.

SEQ ID NO:6 is the deduced amino acid sequence for E-196 polypeptide fragment isolated from tomato, encoded by SEQ ID NO:5.

SEQ ID NO:7 is the nucleotide sequence which codes for H-207 polypeptide fragment isolated from tomato.

5 SEQ ID NO:8 is the amino acid sequence for H-207 polypeptide fragment isolated from tomato, encoded by SEQ ID NO:8.

SEQ ID NO:9 is the nucleotide sequence for the p7-2 fragment isolated from tomato.

10 SEQ ID NO:10 is the nucleotide sequence for the p1-3 fragment isolated from tomato.

SEQ ID NO:11 is the nucleotide sequence for the T6 fragment isolated from tobacco.

SEQ ID NO:12 is the nucleotide sequence for the T1 fragment isolated from tobacco.

15 SEQ ID NO:13 is the nucleotide sequence for the T2 fragment isolated from tobacco.

SEQ ID NO:14 is the nucleotide sequence for the T3 fragment isolated from tobacco.

20 SEQ ID NO:15 is the nucleotide sequence for the PCR1 fragment isolated from tobacco.

SEQ ID NO:16 is the nucleotide sequence for LeMFP1.

SEQ ID NO:17 is the deduced amino acid sequence for LeMFP1, encoded by SEQ ID NO:16.

SEQ ID NO:18 is the nucleotide sequence used as a Southern probe.

25 SEQ ID NO:19 is the nucleotide sequence comprising the cDNA insert in clone src3c.pk004.m1 encoding a soybean MFP1 (GmMFP1).

SEQ ID NO:20 is the deduced amino acid sequence of the nucleotide sequence comprising the cDNA insert in clone src3c.pk004.m1.

30 SEQ ID NO:21 is the nucleotide sequence comprising the cDNA insert in clone p0118.chsab48r encoding a corn MFP1.

SEQ ID NO:22 is the deduced amino acid sequence of the nucleotide sequence comprising the cDNA insert in clone p0118.chsab48r.

SEQ ID NO:23 is the nucleotide sequence comprising the cDNA insert in clone rca1n.pk022.a11 encoding a rice MFP1.

35 SEQ ID NO:24 is the deduced amino acid sequence of the nucleotide sequence comprising the cDNA insert in clone rca1n.pk022.a11.

SEQ ID NO:25 is the nucleotide sequence for PCR primer designed from T3 fragment isolated from tobacco.

SEQ ID NO:26 is the nucleotide sequence for PCR primer designed from T1 fragment isolated from tobacco.

DETAILED DESCRIPTION OF THE INVENTION

The present invention reports the isolation and characterization of cDNAs corresponding to two tobacco MFP1 genes and the isolation and identification of MFP1 EST homologs from corn, soybean and rice. No homologs of MFP1 from tobacco have been described previously. The level of expression of the genes described here can be altered in the plant by methods of cosuppression and overexpression. As they are previously undescribed genes involved in a fundamental cellular mechanism, this can lead to novel developmental phenotypes that might be beneficial for crop growth and development. In addition, if the reduction in expression of one of the genes leads to a growth or developmental defect in the plant, this gene can be used as a novel herbicide target. All isolated proteins can be used as tools to study the plant nuclear matrix, of which no components have been isolated at the molecular level. This can lead to the identification of additional proteins, that can be used as described above. Any related EST sequences can be directly used for the above described applications in crop plants. All of these sequences can be directly used to broaden our understanding of the mechanisms of MAR-matrix interactions and the molecular basis for the described effects on gene expression.

The following definitions are provided for the full understanding of terms and abbreviations used in this specification.

"Polymerase chain reaction" is abbreviated PCR.

"Expressed sequence tag" is abbreviated EST.

"Open reading frame" is abbreviated ORF.

"SDS polyacrylamide gel electrophoresis" is abbreviated SDS-PAGE.

"Amino acid" is abbreviated AA.

"Plaque-forming units" is abbreviated pfus.

" α -Helical" is abbreviated AH.

"Coiled-coil" is abbreviated CC and refers to an amphiphillic α -helical protein structure.

"Hydrophilicity plot" is abbreviated HP.

"Matrix attachment region" is abbreviated MAR. MARs are also known as matrix-associated regions or scaffold-associated (or attachment) regions.

The term "MFP" is an abbreviation for MAR-binding filament-like protein. "MFP1" refers to the MAR-binding filament-like protein having similar characteristics to the protein isolated from tomato as described in Meier et al., *Plant Cell* 8:2105-2115 (1996). "LeMFP1" is the abbreviation for the specific

MFP1 protein isolated from tomato, as set forth in SEQ ID NO:17. "NtMFP1-1" and "NtMFP1-2" are the abbreviations for the first and second MFP1 proteins isolated from tobacco, as set forth in SEQ ID NO:2 and 4 respectively.

5 "GmMFP1" is the abbreviation for the MFP1 protein isolated from soybean, as set forth in SEQ ID NO:20. "ZmMFP1" is the abbreviation for the MFP1 protein isolated from corn, as set forth in SEQ ID NO:22. "OsMFP1" is the abbreviation for the MFP1 protein isolated from rice, as set forth in SEQ ID NO:24. "AtMFP1" is the abbreviation for the MFP1 protein isolated from Arabidopsis, released on (<http://genomewww.standard.edu/Arabidopsis/>).

10 The terms "isolated nucleic acid fragment" or "isolated nucleic acid molecule" refer to a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment or an isolated nucleic acid molecule in the form of a polymer of DNA may be comprised of one or more segments of cDNA,
15 genomic DNA, or synthetic DNA.

The terms "host cell" and "host organism" refer to a cell capable of receiving foreign or heterologous genes and expressing those genes to produce an active gene product. Suitable host cells include microorganisms such as bacteria and fungi, as well as plant cells.

20 The term "fragment" refers to a DNA or amino acid sequence comprising a subsequence of the nucleic acid sequence or protein of the present invention. However, an active fragment of the present invention comprises a sufficient portion of the protein to maintain activity.

The term "substantially similar" refers to nucleic acid fragments wherein
25 changes in one or more nucleotide bases result in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases do not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of
30 the nucleic acid fragments of the present invention such as deletion or insertion of one or more nucleotide bases that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate alteration of gene expression by antisense or co-suppression technology or alteration of the
35 functional properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary sequences.

A "substantial portion" refers to an amino acid or nucleotide sequence which comprises enough of the amino acid sequence of a polypeptide or the

nucleotide sequence of a gene to afford putative identification of that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al., *J. Mol. Biol.* 215:403-410 (1993); see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides (generally 12 bases or longer) may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises enough of the sequence to afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The present specification teaches partial or complete amino acid and nucleotide sequences encoding one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for the purpose known to those skilled in the art. Accordingly, the present invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% identity with the gene to be suppressed. Moreover, alterations in a gene that result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded protein, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the protein molecule would also not be expected to alter the activity of

the protein. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Moreover, the skilled artisan recognizes that substantially similar sequences encompassed by this invention are also defined by their ability to

5 hybridize, under stringent conditions (0.1 x SSC, 0.1% SDS, 65 °C), with the sequences exemplified herein. Preferred substantially similar nucleic acid fragments of the present invention are those nucleic acid fragments whose DNA sequences are 80% identical to the DNA sequence of the nucleic acid fragments reported herein. More preferred nucleic acid fragments are 90% identical to the

10 DNA sequence of the nucleic acid fragments reported herein. Most preferred are nucleic acid fragments that are 95% identical to the DNA sequence of the nucleic acid fragments reported herein.

The term "sequence analysis software" refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid

15 sequences. "Sequence analysis software" may be commercially available or independently developed. Typical sequence analysis software will include but is not limited to the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI), BLASTP, BLASTN, BLASTX (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), and DNASTAR (DNASTAR,

20 Inc. 1228 S. Park St. Madison, WI 53715 USA). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default vales" will mean any set of values or parameters which originally load with the

25 software when first initialized.

The term "percent identity" is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be,

30 as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in: Computational Molecular Biology (Lesk, A. M., ed.) Oxford University Press, New York (1988); Biocomputing: Informatics and Genome Projects (Smith, D. W., ed.) Academic Press, New York (1993);

35 Computer Analysis of Sequence Data, Part I (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); Sequence Analysis in Molecular Biology (von Heinje, G., ed.) Academic Press (1987); and Sequence Analysis Primer (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991).

Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG Pileup program found in the GCG program package, using the Needleman and Wunsch algorithm with their standard default values of gap creation penalty=12 and gap extension penalty=4 (Devereux *et al.*, *Nucleic Acids Res.* 12:387-395 (1984)), BLASTP, BLASTN, and FASTA (Pearson *et al.*, *Proc. Natl. Acad. Sci. USA* 85:2444-2448 (1988)). The BLASTX program is publicly available from NCBI and other sources (BLAST Manual, Altschul *et al.*, Natl. Cent. Biotechnol. Inf., Natl. Library Med. (NCBI NLM) NIH, Bethesda, Md. 20894; Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990); Altschul *et al.*, *IGapped BLAST and PSI-BLAST: a new generation of protein database search programs*, *Nucleic Acids Res.* 25:3389-3402 (1997)). The method to determine percent identity preferred in the present invention is by the method of DNASTAR protein alignment protocol using the Jotun-Hein algorithm (Hein *et al.*, *Methods Enzymol.* 183:626-645 (1990)). Default parameters used for the Jotun-Hein method for alignments are: for multiple alignments, gap penalty=11, gap length penalty=3; for pairwise alignments ktuple=2. As an illustration, for a polynucleotide having a nucleotide sequence with at least 95% identity to a reference nucleotide sequence, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, for a polypeptide having an amino acid sequence having at least 95% "identity" to a reference amino acid sequence, it is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up

to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

“Codon degeneracy” refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the present invention relates to any nucleic acid fragment that encodes all or a substantial portion of present MFP1 proteins as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:20, SEQ ID NO:22 and SEQ ID NO:24. The skilled artisan is well aware of the “codon-bias” exhibited by a specific host cell to use nucleotide codons to specify a given amino acid. Therefore, when synthesizing a gene for improved expression in a host cell, it is desirable to design the gene such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

The term “complementary” is used to describe the relationship between nucleotide bases that are hybridizable to one another. Hence with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine.

A nucleic acid molecule is “hybridizable” to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and solution ionic strength. Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989), particularly Chapter 11 and Table 11.1 therein (entirely incorporated herein by reference). The conditions of temperature and ionic strength determine the “stringency” of the hybridization. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a T_m of 55°, can be used, e.g., 5X SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5X SSC, 0.5% SDS. Moderate stringency hybridization conditions correspond to a higher T_m , e.g., 40% formamide, with 5X or 6X SSC.

Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of
5 complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater
10 than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., *supra*, 9.50-9.51). For hybridizations with shorter nucleic acids, i.e., oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook et al., *supra*, 11.7-11.8). In one embodiment the length for a hybridizable nucleic
15 acid is at least about 10 nucleotides. Preferably a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides; more preferably at least about 20 nucleotides; and most preferably the length is at least 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such
20 as length of the probe.

"Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments which are then enzymatically assembled to construct the entire gene. "Chemically
25 synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on
30 optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determining preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

35 "Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene"

refers to any gene, not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but which is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

"Coding sequence" refers to a DNA sequence that codes for a specific amino acid sequence. "Regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

"Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an "enhancer" is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg, (*Biochemistry of Plants* 15:1-82 (1989)). It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

The "translation leader sequence" refers to a DNA sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner et al., *Mol. Biotech.* 3:225 (1995)).

The "3' non-coding sequences" refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (*Plant Cell* 1:671-680 (1989)).

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA.

"Messenger RNA" (mRNA) refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (U.S. 5,107,065). The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to antisense RNA, ribozyme RNA, or other RNA that is not translated yet has an effect on cellular processes.

The term "operably-linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably-linked with a coding sequence when it affects the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably-linked to regulatory sequences in sense or antisense orientation.

The term "expression" refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the

invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds
5 levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. 5,231,020).

10 "Altered levels" refers to the production of gene product(s) in organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of
15 translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made.
20 "Chloroplast transit sequence" refers to a nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels, J. J., (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can
25 further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the
30 genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al., *Meth. Enzymol.* 143:277 (1987)) and particle-accelerated or "gene gun" transformation technology
35 (Klein et al., *Nature, London* 327:70-73 (1987); U.S. 4,945,050).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook, J., Fritsch,

E.F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

Novel MFP1-binding proteins, have been isolated from tobacco, corn, soybean and rice. Comparison of their random cDNA sequences to the GenBank database using the BLAST and DNASTAR algorithms, well known to those skilled in the art, revealed that these proteins have no significant homologies to other known proteins, other than MFP1 proteins. The nucleotide sequences of the present MFP1 cDNA are provided in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:19, SEQ ID NO:21 and SEQ ID NO:23. Other MFP1 genes and proteins from other plants can now be identified by comparison of random cDNA sequences to the present MFP1 sequences provided herein.

Comparison of the instant MFP1 base deduced amino acid sequences to the only published sequence of this kind (LeMFP1, Meier *et al.*, *Plant Cell* 8:2105-2115 (1996); SEQ ID NO:17 and 18) show a variation of homology of about 39% identity (rice, SEQ ID NO:24) over a length of 107 amino acids to about 77% identity for tobacco (SEQ ID NO:2 and 4) as compared by the Jotun-Hein alignment algorithm (Hein *et al.*, *Methods Enzymol.* 183:626-645 (1990)).

Accordingly preferred polypeptides of the instant invention are those plant proteins which are at least 77% identical to the amino acid sequence as set forth in SEQ ID 17. More preferred amino acid fragments are at least about 80%-90% identical to the sequences herein. Most preferred are nucleic acid fragments that are at least 95% identical to the amino acid fragments reported herein. Similarly, preferred nucleic acid sequences are those encoding MFP1 binding proteins and which are at least 80% identical to the nucleic acid sequences of reported herein. More preferred nucleic acid fragments are at least 90% identical to the sequences herein. Most preferred are nucleic acid fragments that are at least 95% identical to the nucleic acid fragments reported herein.

Similarly preferred polypeptides are those isolated from corn which are at least 40% identical to the polypeptide of SEQ ID NO:17 over a length of about 672 amino acids as compare by the Jotun-Hein alignment algorithm (Hein *et al.*, *supra*). Other preferred polypeptides are those isolated from rice which are at least 39% identical to the polypeptide of SEQ ID NO:17 over a length of about 107 amino acids as compare by the Jotun-Hein alignment algorithm (Hein *et al.*, *supra*). Additionally preferred polypeptides are those isolated from soybean which are at least 46% identical to the polypeptide of SEQ ID NO:17 over a

length of about 388 amino acids as compare by the Jotun-Hein alignment algorithm (Hein et al., *supra*).

5 The nucleic acid fragments of the present invention may be used to isolate cDNAs and genes encoding a homologous MFP1 proteins from the same or other plant species. Isolating homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction (PCR) or ligase chain reaction).

10 For example, other MFP1 genes, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the present nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant using methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the present MFP1 sequences can be designed
15 and synthesized by methods known in the art (Maniatis, *supra*). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primers, DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed and used to
20 amplify a part of or full-length of the present sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the present nucleic acid fragment may
25 be used in PCR protocols to amplify longer nucleic acid fragments encoding homologous MFP1 genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the present nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic
30 acid tracts to the 3' end of the mRNA precursor encoding plant MFP1.

Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al., *Proc. Natl. Acad. Sci., USA* 85:8998 (1988)) to generate cDNAs by using PCR to amplify copies of the region between a single
35 point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the present sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al., *Proc. Natl. Acad. Sci., USA* 86:5673 (1989); Loh

et al., *Science* 243:217 (1989)). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman et al., *Techniques* 1:165 (1989)).

Finally, availability of the present nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the present amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner et al., *Adv. Immunol.* 36:1 (1984); Maniatis, *supra*).

The nucleic acid fragments of the present invention may also be used to create transgenic plants in which the present MFP1 protein is present at higher or lower levels than normal. Alternatively, in some applications, it might be desirable to express the present MFP1 protein in specific plant tissues and/or cell types, or during developmental stages in which they would normally not be encountered.

Overexpression of the present MFP1 may be accomplished by first constructing a chimeric gene in which the MFP1 coding region is operably-linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals must also be provided. The present chimeric genes may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the present chimeric genes can then be constructed. The choice of a plasmid vector depends upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al., *EMBO J.* 4:2411-2418 (1985); De Almeida et al., *Mol. Gen. Genetics* 218:78-86 (1989)), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the MFP1 protein to different cellular compartments or to facilitate their secretion from the cell. The chimeric genes described above may be further modified by the addition of appropriate intracellular or extracellular targeting sequence to their coding regions. These include chloroplast transit peptides (Keegstra et al., *Cell* 56:247-253 (1989), signal sequences that direct proteins to the endoplasmic reticulum (Chrispeels et al., *Ann. Rev. Plant Phys. Plant Mol.* 42:21-53 (1991), and nuclear localization signal (Raikhel et al., *Plant Phys.* 100:1627-1632 (1992). While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of utility may be discovered in the future.

It may also be desirable to reduce or eliminate expression of the MFP1 genes in plants for some applications. In order to accomplish this, chimeric genes designed for antisense or co-suppression of MFP1 can be constructed by linking the genes or gene fragments encoding parts of these enzymes to plant promoter sequences. Thus, chimeric genes designed to express antisense RNA for all or part of MFP1 can be constructed by linking the MFP1 genes or gene fragments in reverse orientation to plant promoter sequences. The co-suppression or antisense chimeric gene constructs could be introduced into plants via well known transformation protocols wherein expression of the corresponding endogenous genes are reduced or eliminated.

The present MFP1 proteins may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to the proteins by methods well known to those skilled in the art. The antibodies would be useful for detecting the present MFP1 protein *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the present MFP1 protein are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the present MFP1. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the present MFP1 protein.

Microbial host cells suitable for the expression of the present MFP1 proteins include any cell capable of expression of the chimeric genes encoding these proteins. Such cells will include both bacteria and fungi including, for example, the yeasts (e.g., *Aspergillus*, *Saccharomyces*, *Pichia*, *Candida*, and *Hansenula*), members of the genus *Bacillus* as well as the enteric bacteria (e.g., *Escherichia*, *Salmonella*, and *Shigella*). Methods for the transformation of such hosts and the expression of foreign proteins are well known in the art and

fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods using PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the present nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function-mutant phenotypes may be identified for the present cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger et al., *Proc. Natl. Acad. Sci. USA* 86:9402 (1989); Koes et al., *Proc. Natl. Acad. Sci. USA* 92:8149 (1995); Bensen et al., *Plant Cell* 7:75 (1995)). The latter approach may be accomplished in two ways. First, short segments of the present nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the MFP1 protein. Alternatively, the present nucleic acid fragment may be used as a hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding a MFP1 protein can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the MFP1 gene product.

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usage and conditions.

35 EXAMPLES

GENERAL METHODS

Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Sambrook et al. (1989), J., Fritsch,

- E. F. and Maniatis, T. Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1989 (hereinafter "Maniatis"); and by T. J. Silhavy, M. L. Bannan, and L. W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory Press, Cold Spring, N.Y. (1984) and by
- 5 Ausubel et al., Current Protocols in Molecular Biology, pub. by Greene Publishing Assoc. and Wiley-Interscience (1987).

Nucleotide and amino acid percent identity and similarity comparisons were made using the DNASTAR suite of programs, applying default parameters unless indicated otherwise.

- 10 The meaning of abbreviations is as follows: "sec" means second(s), "min" means minute(s), "h" means hour(s), "d" means day(s), "μL" means microliter, "mL" means milliliters, "L" means liters, "mM" means millimolar, "M" means molar, "mmol" means millimole(s).

Plant material and growth conditions:

- 15 Tobacco, tomato, soybean, rice, corn, wheat and *Arabidopsis thaliana* were grown in soil in a growth chamber with a 12 h 24 °C light cycle followed by a 12 h, 20 °C dark cycle.

EXAMPLE 1

Isolation Of Total Protein

- 20 Total protein extracts were prepared from leaf tissues. 100 mg aliquots of tissue were ground to a fine powder with mortar and pestle in liquid nitrogen, resuspended in 0.5 mL extraction buffer (62.5 mM Tris-Cl, pH 6.8, 20% glycerol, 4% SDS, and 1.4 M β-mercaptoethanol) and incubated at 70 °C for 10 min. The debris was removed by centrifugation at 15,000 rpm for 10 min at 4 °C. The
- 25 supernatants were removed to new tubes, frozen in liquid nitrogen, and stored at -80 °C.

EXAMPLE 2

Protein Expression, Purification And Antibody Production

- pRSETC-MFP1-EcoRI (containing the coiled-coil domain) and
- 30 pRSETA-MFP1-HincII (containing the DNA binding domain), the expression vectors for E-196 (SEQ ID NO:6) coded by SEQ ID NO:5 and H-207 (SEQ ID NO:8) coded by SEQ ID NO:7 fragments, respectively, have been described previously (Meier *et al.*, *Plant Cell* 8:2105-2115 (1996)). Figure 1 shows a representation of the subfragments E-196 and H-207 that were expressed in
- 35 *Escherichia coli*. Filled bars indicate α-helical regions, open bars indicate hydrophobic domains. The shaded box marks the DNA-binding domain. Numbers indicate the position of the first and last amino acid of each subfragment. Expression of recombinant fusion proteins containing an N-terminal 6-histidine

tag fused to the protein subfragments E-196 (SEQ ID NO:6) and H-207 (SEQ ID NO:8) was induced by isopropyl-D-thiogalactoside in *Escherichia coli* BL21 cells according to the Qiagen protein expression manual (Qiagen, Chatsworth, CA). The amount of fusion protein present in the different total *E. coli* protein extracts was determined by immunoblotting (Maniatis) with a monoclonal antibody directed against the T7 tag (Novagen, Madison, Wisconsin). The expressed proteins were purified by nickel-affinity chromatography (Qiagen, Chatsworth, CA), followed by SDS PAGE. The bands corresponding to the fusion proteins (~1 mg each) were excised from the gel, ground and used to raise two rabbit antisera (a288 against E-196 (SEQ ID NO:6) and aR50 against H-207 (SEQ ID NO:8)). Polyclonal antibodies were produced in rabbits by Eurogentech, Belgium, using the company's standard immunization protocols. The a288 antibody has been described previously (Meier et al., *Plant Cell* 8:2105-2115 (1996)).

EXAMPLE 3

Immuno-detection Of MFP1 Related Proteins

In A Variety Of Higher Plant Species

A 1:3000 dilution of a288 or aR50 antiserum, and a 1:5000 dilution of horseradish peroxidase-coupled anti-rabbit secondary antibody (Amersham, Buckinghamshire, England) were used to perform immunoblot analyses (Maniatis). Enhanced chemiluminescence detection was performed using an ECL detection kit as described by the manufacturer (Amersham Buckinghamshire, England).

a288 and aR50 polyclonal antibodies were then used to detect proteins with antigenic similarity to MFP1 in other plant species (Figure 2). Total protein extracts were prepared from mature leaf tissues of tomato (*Lycopersicon esculentum* L.), tobacco (*Nicotiana tabacum* L.), *Arabidopsis thaliana*, soybean (*Glycine max* L.), rice (*Oryza sativa* L.), wheat (*Triticum aestivum* L.), and corn (*Zea mays* L.) as described above. Equal amounts of total protein, as determined by Coomassie Brilliant Blue staining of a replica protein gel, were probed in immunoblot experiments with aR50 (Figure 2A) and a288 (Figure 2B) polyclonal antibodies. The arrow indicates the position of the MFP1-like proteins of approximately equal size in both panels. The position of molecular weight markers is indicated. aR50 antibody detects a single protein of slightly variable size in all species tested. A second band of higher molecular weight (asterick in Figure 2A) was only occasionally observed in tomato and tobacco extracts (tobacco not shown in Figures 2A-B) and might represent an aggregate of MFP1. In contrast, a288 antibody only detected a protein of about 80 kD in tomato and

tobacco extracts, suggesting that the DNA-binding domain of MFP1 is more highly conserved than the part of the coiled-coil domain present on fragment E-196.

Together, these data indicate that a protein of similar size, containing a related DNA-binding domain, is conserved among higher plant species and that the highest degree of similarity to tomato (LeMFP1) among the plants investigated can be expected in tobacco.

EXAMPLE 4

Cloning and Characterization Of Several Tobacco MFP1 cDNAs

10 Corresponding To Two Tobacco MFP1 Proteins

Example 4 describes the cloning and characterization of two, distinct MFP1 proteins from tobacco.

Isolation Of cDNA By Hybridization

The cDNAs encoding tobacco MFP1 were cloned and characterized. An oligo-dT-primed lambda-ZAP cDNA library made from *Nicotiana tabacum* var. SR1 leaf tissue was purchased from Stratagene (La Jolla, CA). The library was screened in a DNA-hybridization screen according to Maniatis with a 1.6 kb partial cDNA clone representing the 3' 2/3 of the tomato homolog of MFP1, LeMFP1 cDNA (p7 2; SEQ ID NO:9) (Meier et al., *Plant Cell* 8:2105-2115 (1996)) or a 1.0 kb 5' partial LeMFP1 cDNA clone (p1-3; SEQ ID NO:10) (Meier et al., *Plant Cell* 8:2105-2115 (1996)). Hybridization conditions were 5 x Denhards (Maniatis), 5 x SSPE (Maniatis), 5% SDS, 20 µg/mL salmon sperm DNA at 55 °C. Washes were performed at high stringency (0.1 x SSC, 0.1 % SDS at 65 °C). Positive plaques were detected by autoradiography and carried through two subsequent rounds of purification, as described above. *In vivo* excision of positive phage was performed according to the manufacturer's protocol (Stratagene, La Jolla, CA).

Sequencing

In the first screen two positive plaque-forming units (pfus) were detected among approximately 600,000 pfus. After *in vivo* excision, sequence analysis of the two excised cDNAs (T6 (SEQ ID NO:11) and T1 (SEQ ID NO:12)) showed that they represented 1103 bp and 912 bp C-terminal MFP1 sequence, respectively (Figure 3). DNA sequencing was carried out using an ABI Model 377 Sequencer (Perkin Elmer-ABI, Foster City, CA). Sequencing reactions utilized fluorescent sequencing techniques with d-rhodamine and Big Dye terminator chemistry (Perkin Elmer-ABI, Foster City, CA) and were performed according to the standard protocols. The sequence identity between the two 3' fragments T1 (SEQ

ID NO:12) and T6 (SEQ ID NO:11) is 91.5%, suggesting the presence of two MFP1 genes in *Nicotiana tabacum*.

In a second round, the tobacco cDNA library was screened with a 1.0 kb 5' fragment of the LeMFP1 cDNA (p1-3; SEQ ID NO:10) (Meier et al., *Plant Cell* 5 8:2105-2115 (1996)). Two positive pfus were detected among approximately 600,000 pfus. Sequencing of the excised cDNAs (T2 (SEQ ID NO:13) and T3 (SEQ ID NO:14)) showed that they represented partial cDNAs, overlapping with T1 (SEQ ID NO:12) and T6 (SEQ ID NO:11) (Figure 3). Initial sequence analysis of the T2 (SEQ ID NO:13) and T6 (SEQ ID NO:11) cDNAs showed that they 10 shared 445 bp of identical overlapping sequence. It was concluded the T2 (SEQ ID NO:13) and T6 (SEQ ID NO:11) cDNAs represent different portions of the same gene. The overlap of T3 (SEQ ID NO:14) and T1 (SEQ ID NO:12) is only 70 bp, and within this area, there is only a single base pair difference between T6 (SEQ ID NO:11) and T1 (SEQ ID NO:12).

In order to confirm that T3 (SEQ ID NO:14) and T1 (SEQ ID NO:12) 15 were derived from the same gene, PCR primers (SEQ ID NO:25 and SEQ ID NO:26) were designed from the T3 (SEQ ID NO:14) and T1 (SEQ ID NO:12) sequences, that would allow the amplification of a 397 bp fragment, from a *Nicotiana tabacum* lambda ZAP cDNA library, overlapping both cDNAs. PCR 20 reactions were carried out in a Perkin Elmer 9600 thermocycler. (Perkin Elmer, Foster City, CA). The thermocycler was programmed as follows: 2 min 96 °C denaturation cycle, was followed by 30 cycles of 94 °C, 45 sec; 55 °C, 45 sec; 72 °C, 90 sec, and ended with an 8 min 72 °C final extension cycle.

Cloning

25 Using restriction sites added to the primers, the PCR fragment was subsequently cloned into the XbaI/BamHI sites of pSK+ (Stratagene, La Jolla, CA). The sequence of the fragment PCR1 (SEQ ID NO:15) (Figure 3) was found to be 100% identical with both T1 (SEQ ID NO:12) and T3 (SEQ ID NO:14), confirming that these two cDNA fragments are derived from the same gene.

30 Figure 3 shows a schematic structure of the partial tobacco MFP1 cDNAs. T3, T1 and PCR1, shown as open boxes, represent overlapping fragments of the same gene (NtMFP1-1). T2 and T6, shown as filled boxes, represent overlapping fragments of a second gene (NtMFP1-2). The fragment used as a probe for the Southern blot (Figure 5) is indicated.

35 Confirmation Of The Presence Of Two Genes

The divergence between the two tobacco MFP1 cDNAs indicated that they were derived from two different genes. It has been previously shown that a single gene (SEQ ID NO:16) codes for MFP1 (SEQ ID NO:17) in tomato (LeMFP1)

(Meier et al., *Plant Cell* 8:2105-2115 (1996)). Applicants have additionally found that AtMFP1 is a single gene in *Arabidopsis* (data not shown). Based on these findings it was necessary to confirm whether MFP1 is also a single-copy gene in the two diploid progenitors of amphidiploid *Nicotiana tabacum*,

- 5 *N. tomentosiformis* and *N. sylvestris*. In order to confirm this hypothesis the following procedure was applied.

For a Southern blot of *Nicotiana tabacum* genomic DNA, 20 µg aliquots of DNA were digested with various restriction enzymes, run out on 0.8% agarose gel, and were transferred to Immobilon N hydrophobic filters (Millipore, Bedford,
10 MA). Hybridization conditions were essentially as described by Maniatis. The probe (SEQ ID NO:18) was prepared by purification of a 391 bp XhoI/SpeI fragment from the *Nicotiana tabacum* clone T3, as described above. The hybridization temperature was 65 °C. The probe (SEQ ID NO:18), shown in Figure 3, was labelled with ³²P by random prime method according to the
15 manufacturers instructions (BRL, Gaithersburg, MD). Washes were performed at high stringency (0.1 x SSC, 0.1 % SDS at 65 °C).

In the region overlapping the probe, NtMFP1-1 (SEQ ID NO:1) contained a single XbaI site, whereas NtMFP1-2 (SEQ ID NO:3) contains no XbaI site. Neither of the two cDNAs contained an EcoRI site. Figure 5 shows the genomic
20 organization of tobacco MFP1. Tobacco genomic DNA was digested with the indicated restriction enzymes, separated by agarose gel electrophoreses and hybridized in a genomic Southern blot with 391 bp Xho/Spe fragment from the *Nicotiana tabacum* cDNA clone T3 (shown in Figure 3). Abbreviations used are as follows: E, EcoRI; X, XbaI; E/X, EcoRI/XbaI; S, SspI; S/X, SspI/XbaI. The
25 position of DNA size markers is indicated on the right. Two fragments were detected in the lane containing an EcoRI digest (approximately 3.7 kb and 2.7 kb) and three were seen in the lane containing an XbaI digest (approximately 8.0 kb, 7.5 kb and 5.0 kb). In the lane containing the EcoRI/XbaI double digest, the 3.7 kb EcoRI fragment appears to be cleaved by XbaI, leading to two smaller fragments of
30 approximately 1.6 and 0.8 kb. This pattern is consistent with the presence of two genes, one of which contains an XbaI site in the region hybridizing to the probe. In addition, SspI and SspI/XbaI digests were analyzed. Again, one of the two bands detected in the SspI digest is cleaved in the SspI/XbaI double digest. The observed patterns are all consistent with the presence of two genes in the *Nicotiana tabacum*
35 genome, represented by the two isolated cDNAs. These data indicate that, at least in tomato, tobacco and *Arabidopsis*, (data not shown), MFP1 is encoded by a single gene per diploid genome.

In summary, two distinct NtMFP1 cDNAs were isolated from tobacco and named NtMFP1-1 (SEQ ID NO:1) (containing T3 and T1) and NtMFP1-2 (SEQ ID NO:3) (containing T2 and T6). NtMFP1-1 (SEQ ID NO:1) is a full-length cDNA coding for a protein of 722 amino acids (SEQ ID NO:2). NtMFP1-1 (SEQ ID NO:1) and NtMFP1-2 (SEQ ID NO:3) have 77.0% and 78.9% identity to LeMFP1 (SEQ ID NO:16) on DNA level, respectively. The identity between the two tobacco sequences is 91.5%. NtMFP1-1 (SEQ ID NO:2) contains an open reading frame of 721 amino acids. It contains a short 69 bp 5' non-coding region preceding the ATG start codon. NtMFP1-2 (SEQ ID NO:4) contains an open reading frame of 398 amino acids and is not a full-length cDNA.

Table 1 lists the DNASTAR and BLAST comparison of Nt-MFP1-1 and Nt-MFP1-2 with a suit of public databases as well as the literature sequence for tomato (SEQ ID NO:16 and 17) and the MFP1 sequence isolated from Arabidopsis (<http://genomewww.standard.edu/Arabidopsis/>).

15

TABLE I
Comparison of Nt-MFP1-1 and Nt-MFP1-2 Against Known Sequences

| Gene/protein | % Identity* to Tomato | | % Identity* to Arabidopsis | | Similarity Identified | BLAST Algorithm | E value ¹ | Citation |
|--------------|--------------------------|-----------------|-------------------------------|-----------------|--|--------------------|----------------------|---|
| Nt-MFP1-1 | AA ² | NA ³ | AA ² | NA ³ | (Y07861) MFP1 protein [<i>Lycopersicon esculentum</i>] | Xnr | 0.0c | Meier et al., <i>Plant Cell</i> 8:2105-2115 (1996) |
| | 77% | 77% | 42% | 49% | | | | |
| Nt-MFP1-2 | 79% | 78% | 42% | 49% | (Y07861) MFP1 protein [<i>Lycopersicon esculentum</i>] | Xnr | e-147 | Meier et al., <i>Plant Cell</i> 8:2105-2115 (1996) |
| | | | | | | | | |

*Comparison made using DNASTAR MEGALIGN, applying default parameters.

¹Expect value. The Expect value estimates the statistical significance of the match, specifying the number of matches, with a given score, that are expected in a search of a database of this size absolutely by chance.

²AA is the abbreviation for amino acid sequence

³NA is the abbreviation for nucleotide sequence

EXAMPLE 5

Primary And Secondary Structure Analysis Of NtMFP1-1 And NtMFP1-2

Due to the small number of MFP1-like proteins discovered to date, it was advisable to confirm the identity of the present proteins through an analysis of secondary protein structure. Comparisons of the Nt-MFP1-1 and Nt-MFP1-2 proteins were made with the secondary structure of LeMFP1 isolated from tomato and AtMFP1 isolated from *Arabidopsis*.

The *Arabidopsis* genomic DNA sequence was accessed through the *Arabidopsis thaliana* Database (<http://genomewww.standart.edu/Arabidopsis/>).

The deduced protein sequences of the MFP1 proteins were determined and compared using DNASTAR Lasergene software (DNASTAR, Inc., Madison, WI). Figure 4A shows the percent identical amino acids in pairwise comparisons of the four MFP1 proteins.

Based on the amino acid sequence identity NtMFP1-1 (SEQ ID NO:2) and NtMFP1-2 (SEQ ID NO:4) are most closely related. LeMFP1 (SEQ ID NO:17) is more closely related to the two tobacco MFP1s (NtMFP1-1 (SEQ ID NO:2) and NtMFP1-2 (SEQ ID NO:4)) (76% overall sequence identity) than to AtMFP1 (41% overall identity) reflecting the closer relationship of the two solanaeceous species.

Figure 4B shows the hydrophilicity and secondary structure analysis of NtMFP1-1 (SEQ ID NO:2), LeMFP1 (SEQ ID NO:17) and AtMFP1. The secondary structures of the proteins, hydrophilicity, α -helical, and coiled-coil regions were analyzed using DNASTAR PROTEAN software. AH indicates α -helical, CC indicates coiled-coil and HP indicates hydrophilicity plot. The hydrophobic domains are marked with open boxes. Like LeMFP1 (SEQ ID NO:17), NtMFP1 and AtMFP1 contain an extended α -helical, coiled-coil like domain and a shorter N-terminal, non- α -helical region that contains two hydrophobic domains. These structural features are extremely well conserved, despite a relatively low degree of identity on amino acid level in some areas. This is consistent with the more structural conservation of the positioning of polar and non-polar amino acids that is known from other filament-like coiled-coil proteins such as the nuclear lamins (McKeon et al., *Nature* 319:463-468 (1986)). The distance between the first and second hydrophobic domains is very similar in all three proteins (29 AA for tomato, 31 AA for tobacco, and 33 AA for *Arabidopsis* MFP1), indicating a functional relevance of the spacing between the two hydrophobic domains. The length of the N-terminal domain preceding the first hydrophobic domain varies between 56 AA for tomato, 61 AA for tobacco, and 72 AA for *Arabidopsis* MFP1. The common feature of this domain in all three

WO 00/61615

PCT/US00/09723

proteins is a relatively high content of serine and threonine residues (27% to 28%).

EXAMPLE 6

Composition Of cDNA Libraries And Identification

5 Of cDNA Clones From Other Plant Species Encoding Homologs Of MFP1

cDNA libraries representing mRNAs from soybean or corn tissues were prepared. The characteristics of the libraries are described below in Table 2.

Table 2

| cDNA Libraries from Plants | Library | Tissue |
|--|-----------------|--|
| Soybean (<i>Glycine max</i>) | src3c.pk004.m1 | 8 day old root tissue inoculated with eggs of nematode |
| Corn (<i>Zea mays</i>) | p0118.chsab48r | stem tissue, night harvested |
| Rice (<i>Oryza sativa</i> L., Nipponbare) | rca1n.pk022.a11 | callus normalized |

10

Soybean MFP1:

A soybean MFP1 cDNA was identified based on primary and secondary structure analysis. This sequence, from clone src3c.pk004.m1, came from a library prepared from 8 day old root tissue inoculated with eggs of cyst nematode for four days. This sequence contains 1164 base pairs of DNA (SEQ ID NO:19) encoding 388 amino acids (SEQ ID NO:20).

15

Comparison of this partial soybean MFP1 sequence (SEQ ID NO:20) with the sequences from tomato (LeMFP1; SEQ ID NO:17), tobacco (NtMFP1-1; SEQ ID NO:2), and *Arabidopsis* (AtMFP1) shows it to be 46.1, 45.9, and 40.7% identical to these sequences, respectively. In addition, secondary structure analysis of the partial soybean MFP1 (GmMFP1; SEQ ID NO:20) coded by SEQ ID NO:19 shows that it contains an extended α -helical, coiled-coil like domain as do the other MFP1 protein sequences. Results of a Southern blot experiment (not shown) suggest that the soybean MFP1 is encoded by a single copy gene.

20

25 Corn MFP1:

A corn EST sequence was identified as an MFP1 homolog from clone p0118.chsab48r. Secondary structure analysis of the corn MFP1 protein (SEQ ID NO:22) coded by SEQ ID NO:21 shows that it contains an extended α -helical, coiled-coil like domain as well as one of the hydrophobic domains at the N-terminus. Both features are indicative of MFP1 proteins (see Figure 4B).

30

TABLE 3
Comparison of Corn, Rice and Soybean MPF1-1 Sequences Against Known Sequences

| Gene/protein | % Identity* to Tomato | | % Identity* to Arabidopsis | | Similarity Identified | BLAST Algorithm | E value ¹ | Citation |
|--------------|-----------------------|-----------------|----------------------------|-----------------|--|-----------------|----------------------|---|
| | AA ² | NA ³ | AA ² | NA ³ | | | | |
| Corn MPF1 | 40% | 40% | 34% | 48% | (Y07861) MPF1 protein [<i>Lycopersicon esculentum</i>] | Xnr | 2e-94 | Meier et al., <i>Plant Cell</i> 8:2105-2115 (1996) |
| Soybean MPF1 | 46% | 51% | 41% | 48% | (Y07861) MPF1 protein [<i>Lycopersicon esculentum</i>] | Xnr | 5e-76 | Meier et al., <i>Plant Cell</i> 8:2105-2115 (1996) |
| Rice MPF1 | 39% | 43% | 31% | 39% | (Y07861) MPF1 protein [<i>Lycopersicon esculentum</i>] | Xnr | 6e-12 | Meier et al., <i>Plant Cell</i> 8:2105-2115 (1996) |

*Comparison made using DNASTAR MEGALIGN, applying default values

¹Expect value. The Expect value estimates the statistical significance of the match, specifying the number of matches, with a given score, that are expected in a search of a database of this size absolutely by chance.

²AA is the abbreviation for amino acid sequence

³NA is the abbreviation for nucleotide sequence

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|-----------|---|
| (51) International Patent Classification ⁷ : C07K 14/00 | | A2 | (11) International Publication Number: WO 00/61615 |
| | | | (43) International Publication Date: 19 October 2000 (19.10.00) |
| (21) International Application Number: PCT/US00/09723 (22) International Filing Date: 12 April 2000 (12.04.00) (30) Priority Data: 60/128,900 12 April 1999 (12.04.99) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HARDER, Patricia, A. [US/US]; 712 West 34th Street, Wilmington, DE 19802 (US). MEIER, Iris [DE/US]; 4467 Masters Drive, Columbus, OH 43220 (US). (74) Agent: FELTHAM, S., Neil; E.I. Du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). | | | (81) Designated States: BR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i> |
| (54) Title: HOMOLOGS OF MAR-BINDING FILAMENT-LIKE PROTEIN 1 (MFP1) | | | |
| <div style="text-align: center;"><p>MFP1 N — [Diagram of protein structure] — C</p><p>E-196 236 — [Diagram of domain] — 432</p><p>H-207 491 — [Diagram of domain] — 717</p></div> | | | |
| (57) Abstract <p>This invention pertains to nucleic acid fragments encoding proteins that are homologs to the MAR binding filament-like protein 1 (MFP1) from tomato. More specifically, this invention pertains to two tobacco MFP1 genes and MFP1 homologs from corn, soybean, and rice.</p> | | | |

1/4




FIG. 1




FIG. 2A

FIG. 2B

2/4





FIG. 3

3/4

| | N+MFP1-1 | N+MFP1-2 | LeMFP1 | A+MFP1 |
|----------|----------|----------|--------|--------|
| N+MFP1-1 | | 91.5 | 77.0 | 41.9 |
| N+MFP1-2 | | | 78.9 | 42.2 |
| LeMFP1 | | | | 41.5 |

FIG. 4A

FIG. 4B



4/4




FIG. 5

Docket Number
BC1003PCT

DECLARATION and POWER OF ATTORNEY

As a below-named inventor, I hereby declare that.

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

HOMOLOG OF MAR-BINDING FILAMENT-LIKE PROTEIN 1 (MFP1)

the specification of which is attached hereto unless the following box is checked:

☒ was filed on **12 APRIL 2000** as U.S. Application No. _____ or PCT International Application No. _____
PCT/US00/09723 and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

| Application No. | Country | Filing Date | Priority Claimed (Yes/No) |
|-----------------|---------|-------------|---------------------------|
|-----------------|---------|-------------|---------------------------|

I hereby claim the benefit under 35 U.S.C. § 119(c) of any United States Provisional Application(s) listed below.

| U.S. Provisional Application No. | U.S. Filing Date |
|----------------------------------|------------------|
| 60/128,900 | 12 APRIL 1999 |

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International Application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application or PCT International Application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

| Application No. | Filing Date | Status (patented, pending or abandoned) |
|-----------------|-------------|---|
|-----------------|-------------|---|

POWER OF ATTORNEY: I hereby appoint the following attorney(s) and/or agent(s) the power to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

| | |
|--|---|
| Name: S. NEIL FELTHAM | Registration No. 36,506 |
| Send correspondence and direct telephone calls to: S. NEIL FELTHAM | E. I. du Pont de Nemours and Company— Legal—Patents— Wilmington, DE 19898, U.S.A. Tel. No. (302) 992-6460 Fax No. (302) 892-7949 |

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

INVENTOR(S)

| Full Name of Inventor | Last Name | First Name | Middle Name |
|--|---------------------|--------------------------|------------------------|
| HARDER | PATRICIA | | A. |
| Signature (please sign full name): <i>Patricia A. Harder</i> | | | Date: <i>6/1/2000</i> |
| Residence & Citizenship | City | State or Foreign Country | Country of Citizenship |
| WILMINGTON DE | WILMINGTON | DELAWARE | U.S.A. |
| Post Office Address | Post Office Address | City | State or Country |
| 712 WEST 34TH STREET | WILMINGTON | DELAWARE | 19802 |
| Full Name of Inventor | Last Name | First Name | Middle Name |
| MEIER | IRIS | | |
| Signature (please sign full name): <i>Iris Meier</i> | | | Date: <i>6-8-2000</i> |
| Residence & Citizenship | City | State or Foreign Country | Country of Citizenship |
| COLUMBUS OH | COLUMBUS | OHIO | DE |
| Post Office Address | Post Office Address | City | State or Country |
| 4467 MASTERS DRIVE | COLUMBUS | OHIO | 43220 |

☐ Additional Inventors are being named on separately numbered sheets attached hereto.

GENERAL POWER OF ATTORNEY
(Concerning Several International Patent Applications)

The undersigned, Vernon R. Rice, Vice President and Assistant General Counsel of E. I. DU PONT DE NEMOURS AND COMPANY, 1007 Market Street, Wilmington, Delaware 19898 USA ("DuPont"), hereby confirms that the power to sign for DuPont has been granted to various individuals (as set forth in the attached excerpt from DuPont's Patent Board Rules of Procedure (January 1988), Appendix Section III.A.4), including the Chairman, Vice-Chairman, and those individuals who are Assistant Secretaries of the Patent Board. Currently these Assistant Secretaries are:

Roger A. Bowman
Linda J. Davis
John E. Griffiths

Barbara J. Massie
Miriam D. Meconnahey
Deborah A. Meginniss

In addition, the authority to act on behalf of DuPont before the competent International Authorities in connection with any and all international patent applications filed by it with the United States as Receiving Office and to make or receive payments on its behalf is hereby granted to:

| | | | |
|------------------------|--------|--------------------------|--------|
| Beardell, Lori Y. | 34,293 | Kuller, Mark D. | 31,925 |
| Belopolsky, Inna | 43,319 | Krukiel, Charles E. | 27,344 |
| Benjamin, Steven C. | 36,087 | Jarnholm, Arne R. | 30,396 |
| Birch, Linda D. | 38,719 | Langworthy, John A. | 32,255 |
| Bowen, Jr., Alanson G. | 24,027 | Lerman, Bart E. | 31,897 |
| Christenbury, Lynne M. | 30,971 | Levitt, Cary A. | 31,848 |
| Cotreau, William J. | 36,490 | Li, Kening | 44,872 |
| Deitch, Gerald E. | 30,457 | Magee, Thomas H. | 27,355 |
| Deshmukh, Sudhir | 33,677 | Mayer, Nancy S. | 29,190 |
| Dobson, Kevin S. | 40,296 | Medwick, George M. | 27,456 |
| Duffy, Roseanne R. | 33,869 | Morrissey, Bruce W. | 30,663 |
| Edwards, Mark A. | 39,542 | Reynolds, Stephen E. | 37,580 |
| Estrin, Barry | 26,452 | Rizzo, Thomas M. | 41,272 |
| Evans, Craig H. | 31,825 | Santopietro, Lois A. | 36,264 |
| Fair, Tamera L. | 35,867 | Schaeffer, Andrew L. | 33,605 |
| Feltham, S. Neil | 36,506 | Sebree, Chyrrea J. | 45,348 |
| Floyd, Linda Axamethy | 33,692 | Shafer, Robert J. | 24,437 |
| Frank, George A. | 27,636 | Shay, Lucas K. | 34,724 |
| Golian, Andrew G. | 25,293 | Shiple, James E. | 32,003 |
| Gorman, Thomas W. | 31,959 | Siegell, Barbara C. | 30,684 |
| Gould, David J. | 25,338 | Sinnott, Jessica M. | 34,015 |
| Griffiths, John E. | 32,647 | Steinberg, Michael A. | 43,160 |
| Hamby, Jane O. | 32,872 | Steinberg, Thomas W. | 37,013 |
| Hamby, William H. | 31,521 | Stevenson, Robert B. | 26,039 |
| Heiser, David E. | 31,366 | Strickland, Frederick D. | 39,041 |
| Hendrickson, John S. | 30,847 | Tessari, Joseph A. | 32,177 |
| Joung, J. Kenneth | 41,881 | Tulloch, Rebecca W. | 36,297 |
| Katz, Elliott A. | 26,396 | Walker, P. Michael | 32,602 |
| Kelly, Patricia L. | 39,247 | Wang, Chen | 38,650 |
| King, Karen K. | 34,850 | | |

The undersigned ratifies fully all actions already taken by the above-named individuals in accordance with the authority granted hereby.

E. I. DU PONT DE NEMOURS AND COMPANY

By: 

Vernon R. Rice

Vice President and Assistant General Counsel

Date: 9-13-00

SEQUENCE LISTING

<110> E. I. du Pont de Nemours and Company

<120> Homologs of MAR-binding Filament-like protein 1 (MFPL1)

<130> BC1003 PCT

<140>

<141>

<150> 60/128,900

<151> 1999-04-12

<160> 26

<170> Microsoft Office 97

<210> 1

<211> 2168

<212> DNA

<213> Nicotiana tabacum

<400> 1

| | | | | | | |
|------------|------------|-------------|-------------|-------------|-------------|------|
| atggggagtt | cttggtttcc | ccaatctcca | ctctctcatt | ctctcttttc | ttcttcatca | 60 |
| atatcttctt | cccaatttac | acccttgctt | ttttcccca | gaaatgcgca | aaaatgtaaa | 120 |
| aagaaaatgc | cagctatggc | atgtatacac | tcgagagaatc | aaaaggaaag | cgaattctgc | 180 |
| agcagaagaa | cgattctttt | cgtgggtttc | tctgttcttc | cacttctcag | cttgagggca | 240 |
| aatgcttttg | aaggcttgtc | agtagattct | caagtaaaag | cacagccgca | gaaagaggag | 300 |
| acagagcaaa | caatccaagg | aaatgcagag | aatcccttct | tttctctact | taatggactt | 360 |
| ggagtttttg | gttcaggcgt | gcttggttct | ctttatgcct | tggctcgaaa | cgagaaggcc | 420 |
| gtttctgatg | caaccattga | atctatgaaa | aataagctga | aggagaaaga | agccacattc | 480 |
| gtttcatgga | gaagaaattc | cagtctgagc | tgctgaacga | aagggatata | cgaaataatc | 540 |
| aacttaagag | ggcaggcgaa | gaacggcaag | ctctgggtta | ccaattgaat | tcagcaaaga | 600 |
| gtacagtaac | taaccttggt | caggagctgc | aaaaagaaaa | acgaattgct | gaagagctca | 660 |
| tagttcagat | cgagggcctt | caaaataacc | tcatgcagat | gaaggaggat | aagaaaaaat | 720 |
| tgcaggagga | gcttaaagag | aagcttgatt | tgatacaagt | tctgcaagaa | aagataactt | 780 |
| tacttactac | agagatcaaa | gataaagagg | catctcttca | gagtacaacc | tctaaactag | 840 |
| ctgaaaaaga | atcagaggta | gataaattga | gctcaatgta | tcagggaatcc | caggatcagc | 900 |
| tgatgaattt | gacttcagaa | atcaaagaac | ttaaagtcga | agtccagaaa | agagagagag | 960 |
| aactagagtt | gaaacgtgaa | tcagaagaca | accttaaatgt | gcgattaaat | tctttgctcg | 1020 |
| ttgagagaga | tgaatctaaa | aaagagcttg | atgctattca | aaaggaatac | agcgagttca | 1080 |
| agtccatttc | agagaagaaa | gtggcttctg | atgccaaagt | gttgggggaa | caagaaaaga | 1140 |
| gactacacca | gctcgaggaa | caacttgcca | ctgcctcaga | tgaagtacgc | aaaaataatg | 1200 |
| tgctaatacg | tgatctgact | caagaaaaag | aaaacttaag | gagaatgctg | gacgctgagc | 1260 |
| tggaaaacat | aagcaagttg | aagctagagg | tccagggttac | tcaggaaact | cttgagaaat | 1320 |
| ctagaagtga | tgcttctgat | atagcacaac | aactacagca | gtcgaggcat | ctttgctcta | 1380 |
| agcttgaagc | tgaggtttct | aaacttcaga | tggaaattgga | ggaaacaaga | acatcattac | 1440 |
| ggaggaacat | tgatgagaca | aaacgtggtg | caagagctctt | agctgcggag | ctgaccacta | 1500 |
| ctagggagct | tctaaagaaa | acaaatgaag | aaatgcacac | tatgtctcat | gaactagcgg | 1560 |
| ctgttactga | aaattgtgat | aacttacaga | cggagctagt | tgatgtctac | aagaaagcag | 1620 |
| aacgtgctgc | tgatgaactg | aaacaagaaa | agaatattgt | cgtgacactg | gagaaagagc | 1680 |
| taacattttt | ggaggctcaa | attacaagag | agaaagagtc | acggaagaat | ctggaagaag | 1740 |
| agctggaaa | ggctacggaa | tcacttgatg | agatgaaccg | aaatgctttt | gcacttgcaa | 1800 |
| aggagcttga | gcttgctaag | tctcatattt | ctagcctcga | ggatgagaga | gaagtgctcc | 1860 |
| aaaagtctgt | ttctgagcag | aaacaaattt | ctcaagaatc | ccgagaaaac | cttgaagatg | 1920 |
| cccatagcct | ggtaatgaaa | cttggaagg | aacgcgagag | tctggagaag | agagcaaaaga | 1980 |
| aattggaaga | tgaaatggca | tcagcaaaaag | gtgagatttt | gcggctgcgg | acccaagtaa | 2040 |
| attcggtaaa | agctcctggt | aacaatgagg | aaaaagttga | agctggggaa | aaggcagctg | 2100 |
| taacagtga | gagaaccagg | aggaggaaga | ctgctactca | gcctgcttct | cagcaagaaa | 2160 |
| gctcatag | | | | | | 2168 |

<210> 2

<211> 721

WO 00/61615

PCT/US00/09723

<212> PRT

<213> Nicotiana tabacum

<400> 2

| | | | | | | | | | | | | | | | |
|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Met 1 | Gly 2 | Ser | Ser | Cys 5 | Phe | Pro | Gln | Ser | Pro 10 | Leu | Ser | His | Ser | Leu 15 | Phe |
| Ser | Ser | Ser | Ser 20 | Ile | Ser | Ser | Ser | Gln 25 | Phe | Thr | Pro | Leu | Leu 30 | Phe | Ser |
| Pro | Arg | Asn 35 | Ala | Gln | Lys | Cys | Lys 40 | Lys | Lys | Met | Pro | Ala 45 | Met | Ala | Cys |
| Ile | His 50 | Ser | Glu | Asn | Gln | Lys 55 | Glu | Ser | Glu | Phe | Cys 60 | Ser | Arg | Arg | Thr |
| Ile 65 | Leu | Phe | Val | Gly | Phe 70 | Ser | Val | Leu | Pro | Leu 75 | Leu | Ser | Leu | Arg | Ala 80 |
| Asn | Ala | Phe | Glu | Gly 85 | Leu | Ser | Val | Asp | Ser 90 | Gln | Val | Lys | Ala | Gln 95 | Pro |
| Gln | Lys | Glu | Glu 100 | Thr | Glu | Gln | Thr | Ile 105 | Gln | Gly | Asn | Ala | Glu 110 | Asn | Pro |
| Phe | Phe | Ser 115 | Leu | Leu | Asn | Gly | Leu 120 | Gly | Val | Phe | Gly | Ser 125 | Gly | Val | Leu |
| Gly | Ser 130 | Leu | Tyr | Ala | Leu | Ala 135 | Arg | Asn | Glu | Lys | Ala 140 | Val | Ser | Asp | Ala |
| Thr 145 | Ile | Glu | Ser | Met | Lys 150 | Asn | Lys | Leu | Lys | Glu 155 | Lys | Glu | Ala | Thr | Phe 160 |
| Val | Ser | Met | Glu | Lys 165 | Lys | Phe | Gln | Ser | Glu 170 | Leu | Leu | Asn | Glu | Arg 175 | Asp |
| Ile | Arg | Asn 180 | Asn | Gln | Leu | Lys | Arg | Ala 185 | Gly | Glu | Glu | Arg | Gln 190 | Ala | Leu |
| Val | Asn 195 | Gln | Leu | Asn | Ser | Ala | Lys 200 | Ser | Thr | Val | Thr | Asn 205 | Leu | Gly | Gln |
| Glu 210 | Leu | Gln | Lys | Glu | Lys | Arg 215 | Ile | Ala | Glu | Glu | Leu 220 | Ile | Val | Gln | Ile |
| Glu 225 | Gly | Leu | Gln | Asn | Asn 230 | Leu | Met | Gln | Met | Lys 235 | Glu | Asp | Lys | Lys | Lys 240 |
| Leu | Gln | Glu | Glu | Leu 245 | Lys | Glu | Lys | Leu | Asp 250 | Leu | Ile | Gln | Val | Leu 255 | Gln |
| Glu | Lys | Ile 260 | Thr | Leu | Leu | Thr | Thr | Glu 265 | Ile | Lys | Asp | Lys | Glu 270 | Ala | Ser |
| Leu | Gln | Ser 275 | Thr | Thr | Ser | Lys | Leu 280 | Ala | Glu | Lys | Glu | Ser 285 | Glu | Val | Asp |
| Lys | Leu 290 | Ser | Ser | Met | Tyr | Gln 295 | Glu | Ser | Gln | Asp | Gln 300 | Leu | Met | Asn | Leu |
| Thr 305 | Ser | Glu | Ile | Lys | Glu 310 | Leu | Lys | Val | Glu | Val 315 | Gln | Lys | Arg | Glu | Arg 320 |

3

WO 00/61615

PCT/US00/09723

Lys Arg Ala Lys Lys Leu Glu Asp Glu Met Ala Ser Ala Lys Gly Glu
660 665 670

Leu Arg Leu Arg Thr Gln Val Asn Ser Val Lys Ala Pro Val Asn Asn
675 680 685

Glu Glu Lys Val Glu Ala Gly Glu Lys Ala Ala Val Thr Val Lys Arg
690 695 700

Thr Arg Arg Arg Lys Thr Ala Thr Gln Pro Ala Ser Gln Gln Glu Ser
705 710 715 720

Ser

<210> 3
<211> 1199
<212> DNA
<213> Nicotiana tabacum

<400> 3
cgagatgtga atcagaagac aacctgaatg tgcaattaaa ttctttgctc gttgagagag 60
atgaatctaa aaaagagctt gatgctattc aaaaggaata cagcgagttc aagtccattt 120
cagagaagag agtggcttca gatgccaaagc tgttggggga acaagaaaag agactacacc 180
agctcgagga acaacttggt actgccgtaa gtgaagtaag aaaaaataaa gtgctaattg 240
ctaatttgac tcaagcaaaa gaaaacctaa ggagaatgct ggacgctgag ctggaaaatg 300
taagcaagtt gaagctagag gtccagggtta ctccaggaaac tcttgagaaa tcaagaagtg 360
aagcttctga tatagtagaa caactacagc agtcgaggca cttgtgctct aagcttgaag 420
ctgaggtttc taagcttcag atggaattgg aggaaacaag gacattgtta cagaagaaca 480
ttgatgagac aaaacgtggt gcagagttct tagctgcgga gctgaccact actagggagc 540
ttctaaagaa aacaaatgaa gaaatgcaca ccatatccaa tgaactagct gctgttactg 600
aaaatcgtga taacttacag acggagctag ttgatgtcta caagaaagca gaacgtgctg 660
ttaatgaact gaaacaagaa aagaatattg tctgtgacatt ggagaaagag ctaacatttt 720
tggaggctca aattacaaga gagaaagagt cacggaagaa tctggaagaa gagttggaaa 780
gggctacaga atcacttgat gagatgaaca gaaatgcttt tgcacttgca aaggagctgg 840
agctcgctaa ttctcgtatt tctagcctca aagacgagag agaagtgtct caaaagtctg 900
tttctgagca gaagcaaatt tctcaagaag cccgagaaaa ccttgaagat gcccatagcc 960
tgggtgatgaa acttggcaag gaacgcgaga gtctggagaa gagagcaaag aaattggaa 1020
atgaaatggc atcagcaaaa ggtgagattt tgcggttgcg gacacaagta aattcggtaa 1080
aagctcctgt taacaaagag gaaaaagttg aagctgggga aaaggcaaca gtaacagtga 1140
agagaacaac caggaggagg aagactgcta ctctgtctc tcaacaagaa ggctcataa 1199

<210> 4
<211> 398
<212> PRT
<213> Nicotiana tabacum

<400> 4
Arg Cys Glu Ser Glu Asp Asn Leu Asn Val Gln Leu Asn Ser Leu Leu
1 5 10 15

Val Glu Arg Asp Glu Ser Lys Lys Glu Leu Asp Ala Ile Gln Lys Glu
20 25 30

Tyr Ser Glu Phe Lys Ser Ile Ser Glu Lys Arg Val Ala Ser Asp Ala
35 40 45

Lys Leu Leu Gly Glu Gln Glu Lys Arg Leu His Gln Leu Glu Glu Gln
50 55 60

Leu Gly Thr Ala Val Ser Glu Val Arg Lys Asn Lys Val Leu Ile Ala
65 70 75 80

<400> 7

```

gaccactact aaggagcttc taaagaaaac aaatgaagaa atgcacacta tgtcagatga 60
accgtgatag cttacagaca gagctagttg atgtctataa gaaagcagaa catactgcta 120
atgaactgaa acaagaaaag agcattgttg caacactaga agaagagtta aaattttctgg 180
agtctcaaat tacacgagag aaagagttac ggaagagtct ggaagacgag ttagaaaaagg 240
ctacagaatc tcttgatgag attaaccgaa atgtgttggc acttgacagag gagctggagc 300
ttgctacttc tcgtaattct agcctcgaag acgagagaga agtgctccga cagtctgttt 360
ctgagcagaa gcaaatttca caagaagccc aagaaaatct ggaagacgcc catagcctgg 420
tgatgaaact tggcaaggaa cgcgaaagtc ttgagaagag agcaaagaaa ttggaagatg 480
aaatggcagc agcaaaaagg gagattttgc ggctacggag ccaaataaac tcagtaaaaag 540
ctccagtgga ggatgaggaa aaagtgtgtg ctggggaaaaa ggaaaagggtg aaggcaacag 600
taacagcaaa gaaaactacc aggagaagga agagtgtctac tgttaagcaa gaggaaccct 660
ag

```

<210> 8

<211> 226

<212> PRT

<213> Lycopersicon esculentum

<400> 8

```

Thr Thr Thr Lys Glu Leu Leu Lys Lys Thr Asn Glu Glu Met His Thr
  1              5              10              15

Met Ser Asp Glu Leu Val Ala Val Ser Glu Asn Arg Asp Ser Leu Gln
      20              25              30

Thr Glu Leu Val Asp Val Tyr Lys Lys Ala Glu His Thr Ala Asn Glu
      35              40              45

Leu Lys Gln Glu Lys Ser Ile Val Ala Thr Leu Glu Glu Glu Leu Lys
      50              55              60

Phe Leu Glu Ser Gln Ile Thr Arg Glu Lys Glu Leu Arg Lys Ser Leu
      65              70              75              80

Glu Asp Glu Leu Glu Lys Ala Thr Glu Ser Leu Asp Glu Ile Asn Arg
      85              90              95

Asn Val Leu Ala Leu Ala Glu Glu Leu Glu Leu Ala Thr Ser Arg Asn
      100              105              110

Ser Ser Leu Glu Asp Glu Arg Glu Val Leu Arg Gln Ser Val Ser Glu
      115              120              125

Gln Lys Gln Ile Ser Gln Glu Ala Gln Glu Asn Leu Glu Asp Ala His
      130              135              140

Ser Leu Val Met Lys Leu Gly Lys Glu Arg Glu Ser Leu Glu Lys Arg
      145              150              155              160

Ala Lys Lys Leu Glu Asp Glu Met Ala Ala Ala Lys Gly Glu Ile Leu
      165              170              175

Arg Leu Arg Ser Gln Ile Asn Ser Val Lys Ala Pro Val Glu Asp Glu
      180              185              190

Glu Lys Val Val Ala Gly Glu Lys Glu Lys Val Lys Ala Thr Val Thr
      195              200              205

Ala Lys Lys Thr Thr Arg Arg Arg Lys Ser Ala Thr Val Lys Gln Glu
      210              215              220

Glu Pro
225

```

<210> 9
 <211> 1694
 <212> DNA
 <213> *Lycopersicon esculentum*

<400> 9
 gaagagctta aagagaagct tgatttgatt caagttcttg aagaaaagat tacttttgctt 60
 actacagaga tcaaagataa agagggtgagt cttcggagta acacctctaa actagctgaa 120
 aaagaatcgg aggtaaatag tttgagcgat atgtatcaac aatcccagga tcagctgatg 180
 aatttgactt cagagatcaa agaacttaaa gatgaaatcc agaaaagaga gagagaactg 240
 gaggttgaaat gtgtatcaga agacaacctg aatgtgcaat taaattcttt gtcctctcag 300
 agagatgaat ctaaaaaaga gcttcatgct attcaaaagg aatacagtg gttcaagtcc 360
 aattctgatg agaagggtggc ttcagatgag aagctggttg gggacaacaaga gaagagacta 420
 caccagcttg aggaacaact tggcactgcc ttaagtgaag caagtaaaaaa tgaagtgcta 480
 attgctgatc tgactcgaga aaaagaaaaac cttaggagaa tgggtggatgc tgagctggac 540
 aatgtaaaaca agttaaagca agagattgaa gtcactcagg aaagtcttga gaattcaaga 600
 agtgaagttt ctgatataac agtacaacta gagcagttga gggatctttg ctccaaactt 660
 gaagctgagg tttctaaact tcagatggaa ttggaggaaa caagggcatc attacagagg 720
 aacattgatg aaacaaaaa cagttcagag ctcttagctg ctgagttgac cactactaag 780
 gagcttctaa agaaaacaaa tgaagaaatg cacactatgt cagatgaact agtagctgtt 840
 tctgaaaaatc gtgatagctt acagacagag ctgagttgat tctataagaa agcagaacat 900
 actgctaatt aactgaaaca agaaaagagc attgtttgcaa cactagaaga agagttaaaa 960
 tttctggagt ctcaaattac acgagagaaa gagttacgga agagtctgga agacgagtta 1020
 gaaaaggcta cagaatctct tgatgagatt aaccgaaatg tgttggcact tgcagaggag 1080
 ctggagcttg ctacttctcg taattctagc ctcgaagacg agagagaagt gctccgacag 1140
 tctgtttctg agcagaagca aatttcacaa gaagcccaag aaaatctgga agacgcccac 1200
 agcctggtga tgaactttgg caaggaacgc gaaagtcttg agaagagagc aaagaaattg 1260
 gaagatgaaa tggcagcagc aaaagggtgag attttgcggc tacggagcca aataaactca 1320
 gtaaaagctc cagtgaggga tgaggaaaaa gttgttgctg gggaaaagga aaaggtgaag 1380
 gcaacagtaa cagcaaaaga aactaccagg agaaggaga gtgctactgt taagcaagag 1440
 gaaccctagt tggctgtttc tgaatgacat aatcttcttc tttttttgtc ctgactcatt 1500
 tgtttgcaat atttatagag aggccagaat taggacattg ccattggaac aagctgtgta 1560
 ttgtctcttt gagtgatcat ttcccggcga gaagttgcag aaacaaatga ctgatctctt 1620
 gatattcagt caatgttgca gcttactgaa tgaaattatt tgtattgtaa aaaaaaaaaa 1680
 aaaaaaaaaa aaaa 1694

<210> 10
 <211> 1009
 <212> DNA
 <213> *Lycopersicon esculentum*

<400> 10
 taataatggc aacttcttgt tttcctccat tttctgcttc atcttcttca ttatgttctt 60
 cccaattttac accttttgctt tcttgcccaa gaaataccca aatatgtaga aagaagagac 120
 cggttatggc gagtatgcac tcggaaaatc aaaaggaaaag taatgtctgc aacagaagat 180
 cgattctatt tgtgggattc tcagttcttc cacttctcaa tttgagggca agagctctcg 240
 aaggcttgct aacagattct caagcacagc cgcagaaaaga ggaaaccgag caaacaatcc 300
 aaggaagtgc aggaatccc ttcgtttctc tacttaattg acttgggtgtt gttggttcag 360
 gcgtgcttgg ttctctttat gccttggtc gaaatgagaa ggcagtttca gatgcaacca 420
 ttgaatctat gaaaaataag ctgaaggaca aggaagtgc atttgtttca atgaagaagc 480
 aatttgagtc cgaattgctg agcgaaaagg aagatcgaaa taagctaatt aggcgagaag 540
 gtgaagagcg gcaagctttg gttaatcagt taaaatcagc gaagactaca gtaataagcc 600
 ttggctcagga gctgcaaaac gaaaaaaaac ttgctgaaga tctcaaattt gagatcaagg 660
 gccttcaaaa tgacctcatg aatacgaagg aggataagaa gaaattgcag gaagagctta 720
 aagagaagct tgatttgatt caagttcttg aagaaaagat tactttgctt actacagaga 780
 tcaaagataa agaggtgagt cttcggagta acacctctaa actagctgaa aaagaatcgg 840
 agttaaatag tttgagcgat atgtatcaac aatcccagga tcagctgatg aatttgactt 900
 cagagatcaa agaacttaaa gatgaaatcc agaaaagaga gagagaactg gagttgaaat 960
 gtgtatcaga agacaacctg aatgtgcaat taaattcttt gtcctcga 1009

<210> 11
 <211> 1103

<212> DNA
<213> Nicotiana tabacum

<400> 11
cttgagaaat caagaagtga agcttctgat atagtagaac aactacagca gtcgaggcat 60
ctttgctcta agcttgaagc tgaggtttct aagcttcaga tgggaattgga ggaaacaagg 120
acattgtttac agaagaacat tgatgagaca aaacgtggtg cagagttctt agctgcggag 180
ctgaccacta ctagggagct tctaaagaaa acaaatgaag aaatgcacac catatccaat 240
gaactagctg ctgttactga aaatcgatg aacttacaga cggagctagt tgatgtctac 300
aagaaagcag aacgtgctgt taatgaactg aaacaagaaa agaattattgt cgtgacattg 360
gagaaagagc taacattttt ggaggctcaa attacaagag agaaagagtc acggaagaat 420
ctggaagaag agttggaag ggctacagaa tcaattgatg agatgaacag aaatgctttt 480
gcacttgcaa aggagctgga gctcgctaatt tctcgatttt ctacgctcaa agacgagaga 540
gaagtgtctc aaaagtctgt ttctgagcag aagcaatttt ctcaagaagc ccgagaaaaac 600
cttgaagatg cccatagcct ggtgatgaaa cttggcaagg aacgcgagag tctggagaag 660
agagcaaaga aattggaaga tgaaatggca tcagcaaaaag gtgagatttt gcggttgagg 720
acacaagttaa attcggtaaa agctcctgtt aacaaagagg aaaaagttga agctggggaa 780
aaggcaacag taacagtga gagaacaacc aggaggagga agactgctac tctgcttct 840
caacaagaag gctcataatt tgctgtttct gaagtgcac atatccttcc ttttttctt 900
gactcatatt aattgcaacg agggtagatt attggttcat tatataaaac cagaatgagg 960
atattgcctt tgtaagaaac tttctgcaag ctgtattctc agtgagtaaa tttccaggcg 1020
agaagttgcc caaataaatg agatattatt gttgcaagta ccaaatttgg aagggattgt 1080
ttgatatcaa aaaaaaaaaa aaa 1103

<210> 12
<211> 912
<212> DNA
<213> Nicotiana tabacum

<400> 12
atgagacaaa acgtggtgca gagctcttag ctgcggagct gaccactact agggagcttc 60
taaagaaaaa aatgaagaa atgcacacta tgtctcatga actagcggct gttactgaaa 120
attgtgataa cttacagacg gagctagtgt atgtctacaa gaaagcagaa cgtgctgctg 180
atgaactgaa acaagaaaag aatattgtcg tgacactgga gaaagagcta acatttttgg 240
aggctcaaat tacaagagag aaagagtcac ggaagaatct ggaagaagag ctggaaaggg 300
ctacggaatc acttgatgag atgaaccgaa atgcttttgc acttgcaaaag gagcttgagc 360
ttgctaattc tcatatttct agcctcgagg atgagagaga agtgctccaa aagtctgttt 420
ctgagcagaa acaattttct caagaatccc gagaaaacct tgaagatgcc catagcctgg 480
taatgaaact tggcaaggaa cgcgagagtc tggagaagag agcaaaagaaa ttggaagatg 540
aaatggcatc agcaaaagggt gagatttttgc ggctgcggac ccaagtaaat tccgtaaaaag 600
ctcctgttaa caatgaggaa aaagtgaag ctggggaaaaa ggcagctgta acagtgaaga 660
gaaccaggag gaggaagact gctactcagc ctgcttctca gcaagaaaagc tcatagtttg 720
ctgtttctaa gtgacatata tttccttttt gtccttgact caaattgatt gcgacgagaa 780
tagattaatg gtgtattata gagaagccag aattaggata ttgcccttgt aagaaacttc 840
ctgcaagctg tattctcagt gagtgtatat ttccagggtga gaagttgcac aaacaaaaaa 900
aaaaaaaaaa aa 912

<210> 13
<211> 905
<212> DNA
<213> Nicotiana tabacum

<400> 13
cgagatgtga atcagaagac aacctgaatg tgcaattaaa ttctttgctc gttgagagag 60
atgaatctaa aaaagagctt gatgctattc aaaaggaata cagcaggttc aagtcattt 120
cagagaagag agtggcttca gatgccagc tggtggggga acaagaaaag agactacacc 180
agctcgagga acaacttggg actgccgtaa gtgaagtaag aaaaaataaa gtgctaattg 240
ctaatttgac tcaagcaaaa gaaaacctaa ggagaatgct ggacgctgag ctggaaaatg 300
taagcaagtt gaagctagag gtccaggtta ctcaggaaac tcttgagaaa tcaagaagtg 360
aagcttctga tatagtagaa caactacagc agtcgaggca tctttgctct aagcttgaag 420
ctgaggtttc taagcttcag atggaatttg aggaacaag gacattgtta cagaagaaca 480
ttgatagagc aaaacgtggt gcagagctct tagctgcgga gctgaccact actagggagc 540
ttctaaagaa aacaaatgaa gaaatgcaca ccatatccaa tgaactagct gctgttactg 600
aaaatcgtga taacttacag acggagctag ttgatgtcta caagaaagca gaacgtgctg 660

WO 00/61615

PCT/US00/09723

| | | | | | | |
|------------|------------|------------|-------------|------------|------------|-----|
| ttaatgaact | gaaacaagaa | aagaatattg | tctgtgacatt | ggagaaagag | ctaacatttt | 720 |
| tggaggctca | aattacaaga | gagaaagagt | caccgaagaa | tctggaagaa | gagttggaaa | 780 |
| gggctagctc | gcttaagtac | aggagatgga | gaatccaccg | aagaatgaag | tagtggcaga | 840 |
| tcatctgcgt | ccaagcaagt | tacttcacca | acagaaaact | tggatttgta | cctgcctgct | 900 |
| ctccg | | | | | | 905 |

<210> 14
 <211> 1597
 <212> DNA
 <213> *Nicotiana tabacum*

| | | | | | | |
|-------------|------------|-------------|------------|------------|-------------|------|
| <400> 14 | | | | | | |
| cggcctctga | aatcttcttc | tttttatcac | tttcggagtg | gaaatcgga | gaaaccaacc | 60 |
| aactttgtaa | tggggagttc | ttgttttccc | caatctccac | tctctcattc | tctcttttct | 120 |
| tcttcatcaa | tatcttcttc | ccaattttaca | cccttgcttt | tttccccaag | aaatgcgcaa | 180 |
| aaatgtaaaa | agaaaatgcc | agctatggca | tgtatacact | cggagaatca | aaaggaaagc | 240 |
| gaattctgca | gcagaagaac | gattcttttc | gtgggtttct | ctgttcttcc | acttctcagc | 300 |
| ttgagggcaa | atgcttttga | aggcttgctc | gtagattctc | aagtaaaagc | acagccgcag | 360 |
| aaagaggaga | cgagcaacaa | tccaaggaaa | tgcagagaat | cccttctttt | ctctacttaa | 420 |
| tggacttgga | gttttttggt | caggcgtgct | tggttctctt | tatgccttgg | ctcgaaacga | 480 |
| gaaggccgtt | tctgatgcaa | ccattgaatc | tatgaaaaat | aagctgaagg | agaaagaagc | 540 |
| cacattcggt | tcatggagaa | gaaattccag | tctgagctgc | tgaacgaaag | ggatatacga | 600 |
| aataatcaac | ttaagagggc | aggcgaagaa | cggcaagctc | tggttaacca | attgaattca | 660 |
| gcaaaagagta | cagtaactaa | ccttggtcag | gagctgcaaa | aagaaaaacg | aattgctgaa | 720 |
| gagctcatag | ttcagatcga | gggccttcaa | aataacctca | tgcagatgaa | ggaggataag | 780 |
| aaaaaattgc | aggaggagct | taaagagaag | cttgatttga | tacaagttct | gcaagaaaag | 840 |
| ataactttac | ttactacaga | gatcaaagat | aaagaggcat | ctcttcagag | tacaacctct | 900 |
| aaactagctg | aaaaagaatc | agaggtagat | aaattgagct | caatgtatca | ggaatcccag | 960 |
| gatcagctga | tgaatttgac | ttcagaaatc | aaagaactta | aagtcgaagt | ccagaaaaga | 1020 |
| gagagagaac | tagagttgaa | acgtgaatca | gaagacaacc | ttaatgtgct | attaaattct | 1080 |
| ttgctcgttg | agagagatga | atctaaaaaa | gagcttgatg | ctattcaaaa | ggaatacagc | 1140 |
| gagttcaagt | ccatttcaga | gaagaaagtg | gcttctgatg | ccaagctgtt | gggggaacaa | 1200 |
| gaaaagagac | tacaccagct | cgaggaacaa | cttggcactg | cctcagatga | agtacgcaaa | 1260 |
| aataatgtgc | taatcgctga | tctgactcaa | gaaaaagaaa | acttaaggag | aatgctggac | 1320 |
| gctgagctgg | aaaacataag | caagttgaag | ctagaggtcc | aggttactca | ggaaactctt | 1380 |
| gagaaatcta | gaagtgatgc | ttctgatata | gcacaacaac | tacagcagtc | gaggcatctt | 1440 |
| tgctctaagc | ttgaagctga | ggtttctaaa | cttcagatgg | aattggagga | aacaagaaca | 1500 |
| tcattacgga | ggaacattga | tgagacaaaa | cgtggtgcag | agctcttagc | tgccggagctg | 1560 |
| accactacta | gggagcttct | aaagaaaaaa | aaaaaag | | | 1597 |

<210> 15
 <211> 564
 <212> DNA
 <213> *Nicotiana tabacum*

| | | | | | | |
|-------------|-------------|------------|-------------|------------|------------|-----|
| <400> 15 | | | | | | |
| gaggaacaac | ttggcactgc | ctcagatgaa | gtacgcaaaa | ataatgtgct | aatcgctgat | 60 |
| ctgactcaag | aaaaagaaaa | cttaaggaga | atgctggacg | ctgagctgga | aaacataagc | 120 |
| aagttgaagc | tagagggtcca | ggttactcag | gaaactcttg | agaaatctag | aagtgatgct | 180 |
| tctgatatag | cacaacaact | acagcagtcg | aggcatcttt | gctctaagct | tgaagctgag | 240 |
| gtttctaaac | ttcagatgga | attggaggaa | acaagaacat | cattacggag | gaacattgat | 300 |
| gagacaaaaac | gtggtgcaga | gctcttagct | gctggagctga | ccactactag | ggagcttcta | 360 |
| aagaaaaacaa | atgaagaaat | gcacactatg | tctcatgaac | tagcggctgt | tactgaaaat | 420 |
| tgtgataact | tacagacgga | gctagttgat | gtctacaaga | aagcagaacg | tgctgctgat | 480 |
| gaactgaaac | aagaaaagaa | tattgtcgtg | acactggaga | aagagctaac | atttttggag | 540 |
| gctcaaatta | caagagagaa | agag | | | | 564 |

<210> 16
 <211> 2154
 <212> DNA
 <213> *Lycopersicon esculentum*

Arg Asp Leu Cys Ser Lys Leu Glu Ala Glu Val Ser Lys Leu Gln Met
 450 455 460
 Glu Leu Glu Glu Thr Arg Ala Ser Leu Gln Arg Asn Ile Asp Glu Thr
 465 470 475 480
 Lys His Ser Ser Glu Leu Leu Ala Ala Glu Leu Thr Thr Thr Lys Glu
 485 490 495
 Leu Leu Lys Lys Thr Asn Glu Glu Met His Thr Met Ser Asp Glu Leu
 500 505 510
 Val Ala Val Ser Glu Asn Arg Asp Ser Leu Gln Thr Glu Leu Val Asp
 515 520 525
 Val Tyr Lys Lys Ala Glu His Thr Ala Asn Glu Leu Lys Gln Glu Lys
 530 535 540
 Ser Ile Val Ala Thr Leu Glu Glu Glu Leu Lys Phe Leu Glu Ser Gln
 545 550 555 560
 Ile Thr Arg Glu Lys Glu Leu Arg Lys Ser Leu Glu Asp Glu Leu Glu
 565 570 575
 Lys Ala Thr Glu Ser Leu Asp Glu Ile Asn Arg Asn Val Leu Ala Leu
 580 585 590
 Ala Glu Glu Leu Glu Leu Ala Thr Ser Arg Asn Ser Ser Leu Glu Asp
 595 600 605
 Glu Arg Glu Val Leu Arg Gln Ser Val Ser Glu Gln Lys Gln Ile Ser
 610 615 620
 Gln Glu Ala Gln Glu Asn Leu Glu Asp Ala His Ser Leu Val Met Lys
 625 630 635 640
 Leu Gly Lys Glu Arg Glu Ser Leu Glu Lys Arg Ala Lys Lys Leu Glu
 645 650 655
 Asp Glu Met Ala Ala Ala Lys Gly Glu Ile Leu Arg Leu Arg Ser Gln
 660 665 670
 Ile Asn Ser Val Lys Ala Pro Val Glu Asp Glu Glu Lys Val Val Ala
 675 680 685
 Gly Glu Lys Glu Lys Val Lys Ala Thr Val Thr Ala Lys Lys Thr Thr
 690 695 700
 Arg Arg Arg Lys Ser Ala Thr Val Lys Gln Glu Glu Pro
 705 710 715

<210> 18
 <211> 407
 <212> DNA
 <213> Nicotiana tabacum

<400> 18
 tcgaggaaca acttggcact gcctcagatg aagtacgcaa aaataatgtg ctaatcgctg 60
 atctgactca agaaaaagaa aacttaagga gaatgctgga cgctgagctg gaaaacataa 120
 gcaagttgaa gctagaggtc caggttactc aggaaactct tgagaaatct agaagtgatg 180
 cttctgatat agcacaacaa ctacagcagt cgaggcatct ttgctctaag cttgaagctg 240
 aggtttctaa acttcagatg gaattggagg aaacaagaac atcattacgg aggaacattg 300
 atgagacaaa acgtggtgca gagctcttag ctgcggagct gaccactact agggagcttc 360
 taaagaaaaa aaaaaaagga attcctgcag cccgggggat ccactag 407

| | | | | | | |
|------------|-------------|-------------|-------------|-------------|-------------|------|
| ttgacatcac | tgagtgaaga | attcagaaaag | gagaagaaat | tagctgagga | acttagggat | 480 |
| gaaatacaga | gattagagag | tagtatcaca | caagctggca | ttgataatga | tgtgcttgaa | 540 |
| actaaattgg | aagaaaagct | tggtgagatt | aattttttgc | aggaaaaggt | aagtttactc | 600 |
| aaccaagaaa | ttgatgataa | ggagaagcac | atcagggaaac | tcagtgcatac | actttctctcg | 660 |
| aaggaaagta | actacaaaaa | gctgaccgct | ttcacaaatc | aaactaaaaa | gagccttgag | 720 |
| cttgcaaat | ctagagtaca | acaactcgag | gaagaactaa | gtacaactaa | gaacgctctc | 780 |
| gtttctaaga | tatcttctat | tgattcactc | aatgctaaac | ttgaaacctt | gaactctgaa | 840 |
| aagaagaagc | tgacaaaaaa | aataaatgag | ttaatacaag | agtatacaga | cctgaagggt | 900 |
| gcttcagaga | caagggcaag | ccatgattcc | aaactactgt | cagaaagaga | tgatctgata | 960 |
| aaacagcttg | aggaaaaaat | gtctgttgca | ttaactgatt | ctagcaagaa | tcaagaaaaa | 1020 |
| attgttgagt | tgaacaagga | gttgtagtct | accaaaatga | tgctaaagaa | tgaacttaag | 1080 |
| tccatggaag | ctttaaaaaga | ttcaattcga | tcacttgaag | aggctctaaa | gacttcaaga | 1140 |
| agtgaggttt | ccaaactttc | caaggagctt | gaggaggcaa | atgaattgaa | tgaggacctg | 1200 |
| gtatcacaaa | tttctaaact | ccgagaggaa | tccaatgaaa | tgcaagttaga | tctcactaat | 1260 |
| aaactaggag | aggcagaatc | actatctaaa | gctctgtcag | aagatttggtc | ttcagtaaat | 1320 |
| gaaatggttc | agaagggaca | agaagaactc | gaagccacct | ctattgagct | ggcatctatt | 1380 |
| gctgaagctc | gtgacaaact | gaagaaagaa | ttgctggatg | cgtacaagaa | tttgaggtca | 1440 |
| accacacatg | agcttgctga | ggaaagaaaa | attgtgacag | ccttaaaciaa | ggaacttgaa | 1500 |
| gcgttagcga | aacagttgca | ggttgattct | gaagcaagaa | aaagtctcga | atcagacctg | 1560 |
| gaggaggcaa | caaagtcact | agatgaaatg | aacaatagcg | cgctgttact | gtctaaagaa | 1620 |
| cttgagagca | ctcatcttag | gagtgcactg | cttgaaatcg | agaaggaaat | gctacgcaag | 1680 |
| gctctagctg | acacaaacgaa | aatcacacc | gaagctaag | aaaacacaga | ggatgctcag | 1740 |
| aaccttatca | caaggcttga | gacagagaag | gagagctttg | aattgaggtg | tagacatctt | 1800 |
| gaagaggaat | tggcgttagc | aaaaggtgag | atactcgcc | taaggaggca | gattagcaca | 1860 |
| aaagtttctc | agaaacctaa | agcaagagga | ccaccagagg | ccagtgaac | tctgaaggag | 1920 |
| caacctgtga | atgattataa | tcagaagacc | agtgagttg | ttgctggaac | tccacagcct | 1980 |
| gtgaaaagga | ctgtaaggag | aagaaaaggt | agcgcataa | | | 2019 |

```
<210> 22
<211> 672
<212> PRT
<213> Zea mays
```

| <400> 22 | | | | | | | | | | | | | | | | |
|----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| Arg | Thr | Arg | Gly | Pro | Lys | Phe | Glu | Gly | Thr | Lys | Gly | Ile | Ala | Lys | Pro | |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | | |
| Asp | Asn | Thr | Gln | Pro | Glu | Gly | Thr | Gln | Ala | Glu | Thr | Ile | Pro | Glu | Ala | |
| | | | 20 | | | | | 25 | | | | | 30 | | | |
| Arg | Gln | Arg | Glu | Ser | Ser | Leu | Gln | Leu | Val | Gln | Glu | Gln | Pro | Pro | Glu | |
| | | 35 | | | | | 40 | | | | | 45 | | | | |
| Asn | Pro | Leu | Leu | Gly | Phe | Leu | Gly | Ile | Val | Gly | Val | Ala | Ala | Ser | Gly | |
| | 50 | | | | | 55 | | | | | 60 | | | | | |
| Val | Leu | Gly | Gly | Leu | Tyr | Gly | Thr | Ser | Leu | Gln | Glu | Glu | Lys | Ala | Leu | |
| 65 | | | | | 70 | | | | | 75 | | | | | 80 | |
| Gln | Ser | Ile | Val | Ser | Ser | Met | Glu | Ser | Lys | Leu | Ala | Glu | Asn | Glu | Ala | |
| | | | | 85 | | | | | 90 | | | | | 95 | | |
| Ala | Leu | Ser | Leu | Met | Arg | Asp | Asn | Tyr | Glu | Lys | Arg | Leu | Leu | Glu | Gln | |
| | | | 100 | | | | | 105 | | | | | 110 | | | |
| Gln | Ala | Ala | Gln | Lys | Lys | Gln | Ser | Met | Lys | Phe | Gln | Glu | Gln | Glu | Val | |
| | | 115 | | | | | 120 | | | | | 125 | | | | |
| Ser | Leu | Ser | Gly | Gln | Leu | Ala | Ser | Ala | Thr | Lys | Thr | Leu | Thr | Ser | Leu | |
| | 130 | | | | | 135 | | | | | 140 | | | | | |
| Ser | Glu | Glu | Phe | Arg | Lys | Glu | Lys | Lys | Leu | Ala | Glu | Glu | Leu | Arg | Asp | |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Ile | Gln | Arg | Leu | Glu | Ser | Ser | Ile | Thr | Gln | Ala | Gly | Ile | Asp | Asn |
| | | | | 165 | | | | | 170 | | | | | 175 | |
| Asp | Val | Leu | Glu | Thr | Lys | Leu | Glu | Glu | Lys | Leu | Gly | Glu | Ile | Asn | Phe |
| | | | 180 | | | | | 185 | | | | | 190 | | |
| Leu | Gln | Glu | Lys | Val | Ser | Leu | Leu | Asn | Gln | Glu | Ile | Asp | Asp | Lys | Glu |
| | | 195 | | | | | 200 | | | | | 205 | | | |
| Lys | His | Ile | Arg | Glu | Leu | Ser | Ala | Ser | Leu | Ser | Ser | Lys | Glu | Val | Asp |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Tyr | Gln | Lys | Leu | Thr | Ala | Phe | Thr | Asn | Gln | Thr | Lys | Lys | Ser | Leu | Glu |
| 225 | | | | | 230 | | | | | 235 | | | | | 240 |
| Leu | Ala | Asn | Ser | Arg | Val | Gln | Gln | Leu | Glu | Glu | Glu | Leu | Ser | Thr | Thr |
| | | | | 245 | | | | | 250 | | | | | 255 | |
| Lys | Asn | Ala | Leu | Val | Ser | Lys | Ile | Ser | Ser | Ile | Asp | Ser | Leu | Asn | Ala |
| | | | 260 | | | | | 265 | | | | | 270 | | |
| Lys | Leu | Glu | Thr | Leu | Asn | Ser | Glu | Lys | Lys | Lys | Leu | Thr | Lys | Lys | Ile |
| | | 275 | | | | | 280 | | | | | 285 | | | |
| Asn | Glu | Leu | Ile | Gln | Glu | Tyr | Thr | Asp | Leu | Lys | Val | Ala | Ser | Glu | Thr |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Arg | Ala | Ser | His | Asp | Ser | Lys | Leu | Leu | Ser | Glu | Arg | Asp | Asp | Leu | Ile |
| 305 | | | | | 310 | | | | | 315 | | | | | 320 |
| Lys | Gln | Leu | Glu | Glu | Lys | Leu | Ser | Val | Ala | Leu | Thr | Asp | Ser | Ser | Lys |
| | | | | 325 | | | | | 330 | | | | | 335 | |
| Asp | Gln | Glu | Thr | Ile | Val | Glu | Leu | Asn | Lys | Glu | Leu | Asp | Ala | Thr | Lys |
| | | | 340 | | | | | 345 | | | | | 350 | | |
| Met | Met | Leu | Lys | Asn | Glu | Leu | Lys | Ser | Met | Glu | Ala | Leu | Lys | Asp | Ser |
| | | 355 | | | | | 360 | | | | | 365 | | | |
| Ile | Arg | Ser | Ser | Glu | Glu | Ala | Leu | Lys | Thr | Ser | Arg | Ser | Glu | Val | Ser |
| | 370 | | | | | 375 | | | | | 380 | | | | |
| Lys | Leu | Ser | Lys | Glu | Leu | Glu | Glu | Ala | Asn | Glu | Leu | Asn | Glu | Asp | Leu |
| 385 | | | | | 390 | | | | | 395 | | | | | 400 |
| Val | Ser | Gln | Ile | Ser | Lys | Leu | Arg | Glu | Glu | Ser | Asn | Glu | Met | Gln | Val |
| | | | | 405 | | | | | 410 | | | | | 415 | |
| Asp | Leu | Thr | Asn | Lys | Leu | Gly | Glu | Ala | Glu | Ser | Leu | Ser | Lys | Ala | Leu |
| | | | 420 | | | | | 425 | | | | | 430 | | |
| Ser | Glu | Asp | Leu | Ala | Ser | Val | Asn | Glu | Met | Val | Gln | Lys | Gly | Gln | Glu |
| | | 435 | | | | | 440 | | | | | 445 | | | |
| Glu | Leu | Glu | Ala | Thr | Ser | Ile | Glu | Leu | Ala | Ser | Ile | Ala | Glu | Ala | Arg |
| | 450 | | | | | 455 | | | | | 460 | | | | |
| Asp | Asn | Leu | Lys | Lys | Glu | Leu | Leu | Asp | Ala | Tyr | Lys | Asn | Leu | Glu | Ser |
| 465 | | | | | 470 | | | | | 475 | | | | | 480 |
| Thr | Thr | His | Glu | Leu | Val | Glu | Glu | Arg | Lys | Ile | Val | Thr | Ala | Leu | Asn |
| | | | | 485 | | | | | 490 | | | | | 495 | |

Lys Glu Leu Glu Ala Leu Ala Lys Gln Leu Gln Val Asp Ser Glu Ala
 500 505 510
 Arg Lys Ser Leu Glu Ser Asp Leu Glu Glu Ala Thr Lys Ser Leu Asp
 515 520 525
 Glu Met Asn Asn Ser Ala Leu Leu Leu Ser Lys Glu Leu Glu Ser Thr
 530 535 540
 His Ser Arg Ser Ala Thr Leu Glu Ser Glu Lys Glu Met Leu Arg Lys
 545 550 555 560
 Ala Leu Ala Glu Gln Thr Lys Ile Thr Thr Glu Ala Lys Glu Asn Thr
 565 570 575
 Glu Asp Ala Gln Asn Leu Ile Thr Arg Leu Glu Thr Glu Lys Glu Ser
 580 585 590
 Phe Glu Leu Arg Cys Arg His Leu Glu Glu Glu Leu Ala Leu Ala Lys
 595 600 605
 Gly Glu Ile Leu Arg Leu Arg Arg Gln Ile Ser Thr Asn Ser Ser Gln
 610 615 620
 Lys Pro Arg Ala Arg Gly Pro Pro Glu Ala Ser Glu Thr Leu Lys Glu
 625 630 635 640
 Gln Pro Val Asn Asp Tyr Asn Gln Lys Thr Ser Gly Val Val Ala Gly
 645 650 655
 Thr Pro Gln Pro Val Lys Arg Thr Val Arg Arg Arg Lys Gly Gly Ala
 660 665 670

<210> 23
 <211> 322
 <212> DNA
 <213> Oryza sativa

<220>
 <223> n= g, a, c or t

<400> 23
 gagagaaact agttctagga aggacactct tgaagcagag aaaaaaatgt tatcaaaggc 60
 tcttgctgag caacagaaga tcacaactga agctcatgaa aacactgagg atgctcagaa 120
 tcttatctct aggccttcaga ctgagaagga gagttttgaa atgagggcta gacatcttga 180
 agaggagttg gcgtagcaaa aggggtgagat attgcgccta agaaggcaga ttagtacaag 240
 cagatcacag aaagcaaaaa ctcttccaaa cacaaatgca tctccagagg tcagtcaggc 300
 tccangacga gcaggctgtg aa 322

<210> 24
 <211> 107
 <212> PRT
 <213> Oryza alta

<220>
 <223> X= G or R

<400> 24
 Arg Glu Thr Ser Ser Arg Lys Asp Thr Leu Glu Ala Glu Lys Lys Met
 1 5 10 15
 Leu Ser Lys Ala Leu Ala Glu Gln Gln Lys Ile Thr Thr Glu Ala His
 20 25 30

